

RESEARCH ARTICLE

Derivation of physiological inhalation rates in children, adults, and elderly based on nighttime and daytime respiratory parameters

Pierre Brochu¹, Jules Brodeur², and Kannan Krishnan²

¹Ministère du Développement durable, de l'Environnement et des Parcs, Direction du suivi et de l'état de l'environnement, Service des avis et expertises scientifiques, gouvernement du Québec, édifice Marie-Guyart, Québec, QC, Canada, and

²Département de santé environnementale et santé au travail, Faculté de médecine, Université de Montréal, succursale Centre-Ville, Montréal, QC, Canada

Abstract

The methodology developed in our previous studies (Brochu *et al.* 2006a–c) for the determination of physiological daily inhalation rates was improved by integrating into the calculation process, both nighttime and daytime respiratory parameters, namely oxygen uptake factors (H) and ventilatory equivalents (VQ). H values during fasting (0.2057 ± 0.0018 L of O_2 /kcal; mean \pm SD) and postprandial phases (0.2059 ± 0.0019 L of O_2 /kcal) as well as VQ values for subjects at rest (27.4 ± 4.8 to 32.2 ± 3.1 , unitless) and during the aggregate daytime activities (29.9 ± 4.2 to 33.7 ± 7.2) were determined and combined with published doubly labeled water measurements for the calculation of daily inhalation rates in normal-weight males and females aged 0.22–96 years ($n = 1235$). Depending upon the unit value chosen, the highest 99th percentiles for inhalation data were found in males aged 35 to <45 years (35.40 m³/day), 2.6 to <6 months (1.138 m³/kg-day), and 10 to <16.5 years (22.29 m³/m²-day). Means and percentiles expressed in m³/kg-day as well as in m³/m²-day suggest generally higher intakes of air pollutants in children than in adults and in males than in females (in μ g/kg-day or μ g/m²-day) for identical exposure concentrations and conditions. For instance, means in boys aged 2.6 to <6 months of 10.99 ± 3.50 m³/m²-day and 0.572 ± 0.191 m³/kg-day are 1.3- and 2.5-folds higher, respectively, than those in adult males 65–96 years old (8.42 ± 2.13 m³/m²-day, 0.225 ± 0.059 m³/kg-day).

Keywords: Daily inhalation rates, oxygen uptake factor, ventilatory equivalent, doubly labeled water, health risk assessment

Introduction

Accurate values for daily inhalation rates in humans are required for health risk assessment and management of air pollutants (Health Canada 1996; van Engelen and Prud'homme de Lodder 2007) especially for the young and aged, who are thought to be more susceptible than adults to the adverse health effects of airborne chemicals (Braun-Fahrlander *et al.* 1997; Tolbert *et al.* 2000; Liu *et al.* 2003; Yang *et al.* 2003).

Estimates of daily inhalation rates in humans have been greatly improved with the use of the energy expenditure approach of Layton (1993). This approach has been formulated in a basic equation comprising the following

terms (Equation 1): E (mean energy expenditure required for a given activity level expressed as kcal/min), H (oxygen uptake factor expressed as L of oxygen consumed/kcal expended), and VQ (ventilatory equivalent ratio of the minute ventilation rate (VE) to the oxygen consumption rate (VO_2), unitless). Nevertheless, the procedures developed by Layton (1993) to estimate E values are not free from biases and were showed to generate errors of daily inhalation estimates ranging from –36% to +60% (Brochu *et al.* 2006c). The difficulty in achieving accurate estimations of E values has been addressed by Brochu *et al.* (2006a, b) with the use of total daily energy expenditures (TDEE) that are measured from the doubly labeled

Address for Correspondence: Pierre Brochu, Ministère du Développement durable, de l'Environnement et des Parcs, Direction du suivi et de l'état de l'environnement, Service des avis et expertises scientifiques, gouvernement du Québec, édifice Marie-Guyart, 7^e étage, 675, boulevard René-Lévesque Est, Québec, QC, G1R 5V7, Canada. E-mail: pierre.brochu@mddep.gouv.qc.ca

(Received 13 July 2010; revised 16 November 2010; accepted 24 November 2010)

Abbreviations

α	data for the aggregate daytime activities of subjects	RER	VCO_2/VO_2 ratio, more properly known as the respiratory exchange ratio
β	data for subjects under resting conditions	Sld	sleep duration
BEE	basal energy expenditure (BMR expressed on a 24-h basis) BMI body mass index	SMR	sleeping metabolic rate
BMR	basal metabolic rate (punctual measurement)	STPD	standard temperature and pressure, dry air
BSA	body surface area	TDEE	total daily energy expenditure
BTPS	body temperature pressure saturation	VCO_2	carbon dioxide production rate
DLW	doubly labeled water	VE	minute ventilation rate
E	minute energy expenditure rate	VO_2	oxygen consumption rate (also known as the oxygen uptake)
ECG	stored daily energy cost for growth	VQ	ventilatory equivalent for VO_2 (VE at BTPS/ VO_2 at STPD)
H	oxygen uptake factor, volume of oxygen (at STPD) consumed to produce 1 kcal of energy expended		

water (DLW) method (Bluck 2008). Values for TDEE systematically encompass voluntary and involuntary energy expended by humans during real-life situations in their normal surroundings each minute of the day, 24 h per day, on a daily basis for 7–21 days (IDECG 1990).

The precise values of the two other parameters in the equation of Layton (i.e. H and VQ) are still a matter for discussion. A postprandial H value of 0.21 L of O_2 /kcal has been first calculated for Americans by Layton (1993) and later confirmed for Canadians by Brochu *et al.* (2006a). However, a critical analysis of possible fluctuations in the postprandial H value as a function of age, sex, and typical dietary intakes in various countries has not been performed. Moreover, the variation of H values during nighttime sleep in fasting subjects has never been taken into account in the calculation process of daily inhalation rates. Similarly, VQ values have been shown to vary from 34.2 to 36.8 in pregnant and lactating women in Brochu *et al.* (2006b) compared with the constant value of 27 reported in Layton (1993). However, the accurate variation of VQ values in non-gestational and lactating individuals as a function of age has not yet been reliably characterized.

The present article is therefore intended to improve the methodology developed previously by Brochu *et al.* (2006a–c) for a scientifically sound determination of daily inhalation rates in free-living individuals based on DLW measurements. The overall approach involved the determination and integration of the means and standard deviations for E , H , and VQ for nighttime sleep (fasting phase) and daytime activities (postprandial phase) into the calculation process of physiological daily inhalation rates in normal-weight individuals.

Methodology

Study design

Means and standard deviations (SD) for H and VQ were determined initially and then used subsequently in the second part, with those of E and sleep durations (Sld), for the calculation of the physiological daily inhalation rates. Data for athletes and explorers were excluded from the calculation process of the latter values. Daily inhalation values were expressed as absolute values (m^3/day),

as well as relative values to the body weight ($\text{m}^3/\text{kg}\cdot\text{day}$) and body surface area (BSA; $\text{m}^3/\text{kg}\cdot\text{m}^2$). Normal-weight individuals were defined according to the following body mass index (BMI) cutoffs: from the 3rd to 97th percentiles for children under 3 years old, the 85th percentile or below for children aged 3–19 years, and from 18.5 to 24.5 kg/m^2 for adults over 19 up to 96 years (IOM 2002). Infants, toddlers, children, and teenagers are hereafter collectively referred to children.

Values for E were determined by using individual DLW measurements taken from the database reported in IOM (2002) for healthy normal-weight males and females aged 2.6–96 years ($n=1235$). These values, which are systematically measured with the DLW method, include subject-specific information on body weight, height, BMI value, basal energy expenditure (BEE), and TDEE values. Values for BEE were measured by indirect calorimetry (Ferrannini 1988; Bursztein *et al.* 1989), whereas those for TDEE were obtained by mass spectrometric monitoring of disappearance rates of oral doses of water isotopes usually monitored in the urine (IDECG 1990). Values for E during nighttime sleep were calculated by using BEE values. Those during the aggregate daytime activities are the result of subtracting BEE from TDEE values.

An exhaustive compilation and a critical analysis of published data in healthy subjects were performed in order to select appropriate parameters for the determination of H values during postprandial and fasting phases (i.e. typical diets found in various countries, respiratory gas-exchange measurements of oxygen and carbon dioxide) and VQ values under resting conditions and for the aggregate daytime activities (i.e. simultaneous measurements of minute ventilation and VO_2) (appendix).

Food recall surveys (i.e. retrospective method) or weighed dietary records (i.e. prospective method using household measures or collection of duplicate diets) are used to describe dietary intakes in subjects (Torun *et al.* 1996). Experimental procedures used for measurements of VO_2 , carbon dioxide production (VCO_2), and VE are specified in each publication. However, VO_2 and VCO_2 values are often measured using paramagnetic O_2 and infrared CO_2 analyzers, respectively (Skoog *et al.* 2006). Values for VE are generally measured by spirometry and sometime by

pneumotacography (Mason *et al.* 2005). Sld are recorded day-by-day on questionnaires by survey respondents for extensive periods of time (usually longer than a year) including complementary data, as those regarding work conditions, physical activities, diets, as well as health and socioeconomic variables (e.g. Bjorvatn *et al.* 2007).

VO₂β and VO₂α: criteria for data selection for H and VQ calculations

Published sets of measurements for VE, VO₂, and VCO₂, VO₂ values measured in healthy subjects at rest or while performing various activities at about the sea level, when breathing an oxygen concentration of 21%, were ranked per age groups. Then, only those measured in subjects with experimental VO₂ demands within the span of VO₂ values for resting conditions (referred to as β) or the aggregate daytime activities (referred to as α) were included in the present study. Values for VO₂β and VO₂α were calculated by using BEE and TDEE values reported in the database of the IOM (2002) for healthy normal-weight individuals (age = 2.6 months–96 years; *n* = 1235). According to Layton (1993), VE (L/min) is expressed as a function of *H* (L of O₂/kcal), *E* (kcal/min), and VQ (i.e. VE/VO₂ ratio, unitless) values as follows:

$$VE = E \times H \times VQ \quad (1)$$

Hence,

$$VO_2 = E \times H \quad (2)$$

where *H* is the volume of oxygen consumed at standard temperature and pressure, dry air (STPD) to produce 1 kcal of energy expended, and VQ is the ratio of the VE value at body temperature and saturated with water vapor (BTPS) to the VO₂ value at STPD.

Therefore, values for minute energy expenditure rates (Eβ and Eα in kcal/min) as well as VO₂β and VO₂α (L/min) were expressed in terms of BEE and TDEE values (kcal/day) as well as the daily energy costs for growth (ECG, in kcal/day) and Sld (in h/day) by using the following equations:

$$E\beta = \left[\frac{BEE + ECG}{1440} \right] \quad (3)$$

$$E\alpha = \left[\frac{TDEE - BEE}{(24 - Sld) \times 60} \right] + \left[\frac{BEE + ECG}{1440} \right] \quad (4)$$

$$VO_{2\beta} = \left[\frac{BEE + ECG}{1440} \right] \times H \quad (5)$$

$$VO_{2\alpha} = \left[\frac{(TDEE - BEE)}{(24 - Sld) \times 60} + \frac{(BEE + ECG)}{1440} \right] \times H \quad (6)$$

where 1440 and 60 are the conversion factors from days to minutes and hours to minutes, respectively, and 24 is the number of hours in a day.

Values for ECG were added to BEE values in order to take into account the energy demands required during the growth process from birth up to 18 years of age for females and 24 years old for males (Brochu *et al.* 2006a). The BEE value corresponds to the basal metabolic rate (BMR) expressed during a 24-h period. The BMR value is defined as the sum of the total energy expenditure required to maintain the minimal tissue cellular activity in order to sustain vital functions, notably blood circulation, respiration, gastrointestinal, and renal processes (Guyton 1991). BMR values are measured under standard conditions in a comfortably warm room, with subject lying at complete rest in thermoneutral conditions and having fasted for 12–13 h. Respiratory gas-exchange rates are measured for subjects 40 min immediately after waking (e.g. Butte *et al.* 2004). The postprandial *H* value of 0.21 L of O₂/kcal used by Layton (1993) and Brochu *et al.* (2006a–c) is in accordance with that calculated in the present study (0.207 L of O₂/kcal) by using VO₂ and BMR values per unit of organ weight established by Malcom and Hollyday (1971). Values for VO₂ per unit of tissue weight (3.7 to 123.8 L of O₂/kg-day) for brain, liver, heart, kidneys, and muscles reported in Malcom and Hollyday (1971) for adults correspond to a mean *H* value of 0.207 L of O₂/kcal for these five organs when divided by their respective BMR (17.6–606 kcal/kg of organ per day). Consequently, a *H* value of 0.21 L of O₂/kcal was used for the calculation of the lower and upper limits of VO₂β and VO₂α for the different age groups in this study (Tables 1 and 2).

H values

Variations of the postprandial *H* value (referred to as *H_p* value) as a function of age, sex, and country were calculated based on typical dietary intake contributions found in 17 countries. This is done by taking into account absorption rates of ingested protein, fat, and carbohydrates (92%, 95%, and 98%, respectively) through the gastrointestinal tract (Guyton 1991) and considering that the oxidation of 1 g each of these nutrients consumes 0.97, 0.83, and 2.0 L of O₂ and yields 4.5, 9.5, and 4.2 kcal of energy, respectively (McLean and Tobin 1987; Layton 1993; Brochu *et al.* 2006a). Values for *H_p* and *H* for fasting subjects (referred to as *H_F* value) were also calculated by using values for VO₂ and VCO₂, or alternatively using VO₂ and respiratory exchange ratios (i.e. VCO₂/VO₂, known as the RER value) simultaneously measured by indirect calorimetry at STPD in the same subjects. Then, values for VO₂ and VCO₂ (L/min) were converted into minute energy expenditure rate (*E*, kcal/min) and *H* (L/kcal) by using the following equations (Weir 1949):

$$E = 3.941 \times VO_2 + 1.106 \times VCO_2 \quad (7)$$

$$H = VO_2 \times (3.941 \times VO_2 + 1.106 \times VCO_2)^{-1} \quad (8)$$

The combustion of carbohydrates, protein, and fat from ingested food requires 0.199, 0.212, and 0.221 L of O₂ per kcal of energy expended, respectively (McLean and

Table 1. Anthropometric, energetic measurements, and oxygen consumption rates in healthy normal-weight males and females aged 2.6 months to <10 years.

2.0 months to <10 years.															
Gender and age group (years)		Body weight (kg) ^a		Body surface area (m ²)		Energetic measurement (kcal/day)						Oxygen consumption rate (L/min)			
		Mean ± SD		D		BEE ^b		ECG ^c		TDEE ^d		VO ₂ β ^e		VO ₂ α ^e	
						Mean ± SD	D	Mean ± SD	D	Mean ± SD	D	Min	Max	Min	Max
Males															
0.22 to <0.5	28	6.6±1.0	L	0.34±0.03	L	387±64	L	121±42	L	492±125	L	0.06	0.09	0.06	0.19
0.5 to <1	37	8.8±1.1	L	0.42±0.03	L	532±63	N	40±9	L	722±123	L	0.07	0.10	0.08	0.24
1 to <2	34	10.7±1.1	N	0.49±0.04	N	668±71	N	22±4	L	890±145	L	0.07	0.12	0.11	0.28
2 to <5	25	15.3±3.4	N	0.64±0.10	N	846±153	N	17±4	L	1176±274	L	0.09	0.16	0.13	0.35
5 to <7	96	19.8±2.1	L	0.79±0.06	L	1012±91	N	41±6	L	1398±192	L	0.12	0.20	0.18	0.45
7 to <10	28	26.8±4.2	L	0.98±0.10	L	1129±116	N	51±10	L	1722±322	L	0.13	0.21	0.18	0.55
Females															
0.22 to <0.5	49	6.6±0.9	L	0.34±0.03	L	374±53	L	117±42	L	471±102	L	0.06	0.09	0.07	0.20
0.5 to <1	63	8.5±1.0	L	0.41±0.03	L	506±67	L	38±7	L	661±121	N	0.06	0.10	0.07	0.24
1 to <2	61	10.6±1.4	L	0.49±0.04	L	630±85	L	18±3	L	844±160	N	0.07	0.13	0.09	0.28
2 to <5	36	14.4±3.0	L	0.62±0.09	L	776±132	N	19±4	L	1083±219	L	0.08	0.16	0.11	0.34
5 to <7	102	19.7±2.3	L	0.79±0.06	L	943±75	N	34±5	L	1332±184	L	0.12	0.17	0.16	0.39
7 to <10	140	27.3±3.6	L	0.99±0.08	L	1079±86	N	42±7	L	1660±265	L	0.13	0.20	0.19	0.51

N = normal; L = lognormal. n = number of individuals; SD = standard deviation.

^aNormal-weight for children aged 2.6 months to <3 years with body mass index (BMIs) between the 3rd and the 97th percentiles and those aged 4 to <10 years with BMIs corresponding to the 85th percentile or below (IOM 2002).

^bBEE = basal energy expenditure (i.e. basal metabolic rate expressed on a 24-h basis) measured by indirect calorimetry (IOM 2002).

^cECG = stored daily energy cost for growth (Brochu *et al.* 2006a).

^dTDEE = total daily energy expenditure. TDEEs were based on ²H₂O and H₂¹⁸O disappearance rates from urine monitored by gas-isotope ratio mass spectrometry during 7- to 21-day periods for free-living individuals (IOM 2002).

^eVO₂β and VO₂α = oxygen consumption rates for individuals at rest and during aggregate daytime activities, respectively, used for the selection of data for the calculation of H and VQ values. D = best fit distribution (i.e. lognormal or normal) according to the Anderson-Darling goodness-of-fit test performed on individual data for each age group.

Tobin 1987). During the fasting phase, 0.198, 0.200, 0.210, 0.211, and 0.214 L of O₂/kcal are required for the combustion of glycogen, glucose, 3-hydroxybutyric acid, acetoacetic acid, and triacylglycerol, respectively (Elia 1997). Consequently, a minimum of 0.199 L of O₂/kcal and maximum of 0.221 L of O₂/kcal for H_p values (McLean and Tobin 1987), as well as minimal and maximal H_p values of 0.198 and 0.214 L of O₂/kcal, respectively (Elia 1997), were used into the calculation process of physiological daily inhalation rates.

VQ values

Values for VQ were calculated by dividing VE by VO₂ values simultaneously measured for the same subjects at BTPS and STPD, respectively. Voluntary and involuntary activities during daytime are performed by individuals in the sitting or standing position. Therefore, VQα values were calculated exclusively by using published VEα and VO₂α measured while subjects were in the upright position. The data for subjects in the supine position were insufficient to calculate VQβ values. However, only slightly higher energy expenditure is required when subjects, during resting conditions, change from a supine to an upright position, which consequently increase VO₂, VCO₂, VE values by about the same extent (e.g. Donevan *et al.* 1962; Damato *et al.* 1966). Conversely, lower BMR values observed in normal-weight subjects during profound sleep (e.g. Ravussin *et al.* 1985; Garby *et al.*

1987) slightly reduce VE and VO₂ demands as well (e.g. Colrain *et al.* 1987). These fluctuations of VO₂ demands combined with the change of VE and VO₂ values always remain within the span of VO₂β. Therefore, VQβ values were calculated by using sets of VEβ and VO₂β values measured in subjects in the upright position. Such VQβ values can be used to characterize VQ values for subjects during resting conditions in the upright or supine position as well as during nighttime sleep.

Published sets of VE and VO₂ values were found for individuals aged <1 year and for those from 4 to 91 years in the supine and upright positions, respectively. No data were available for children from 1 to <4 years of age. Thus, VQ values for the latter aged group were assumed to be the same as those for children aged 1 to <10 years. Mean VQβ values of 30.2 ± 7.6 and 30.8 ± 0.9 calculated for nonsedated children aged 2 h to 1.4 months (Cook *et al.* 1955; Stahlman and Meece 1957; Nelson *et al.* 1962; n = 131) and 4 to <10 years, respectively (Robinson 1938; Inbar *et al.* 1981; n = 35), are within the same order of magnitude, and both appear to be slightly higher than the value of 27.0 ± 4.3 based on data reported in Lees *et al.* (1967) for sedated children aged 0.5–8.5 months (n = 26). Therefore, the former value (i.e. 30.2 ± 7.6) was used to characterize VQβ in children aged 2.6 months to <1 year rather than the latter (i.e. 27.0 ± 4.3). The VQα value for children aged <1 year old was assumed to be the same as the VQβ value since such children have limited

Table 2. Anthropometric, energetic measurements, and oxygen consumption rates in healthy normal-weight males and females aged 10–96 years.

Gender and age group (years)		Body weight (kg) ^a		Body surface area (m ²)		Energetic measurement (kcal/day)						Oxygen consumption rate (L/min)			
		Mean ± SD	D	Mean ± SD	D	BEE ^b		ECG ^c		TDEE ^d		VO ₂ β ^e		VO ₂ α ^e	
						Mean ± SD	D	Mean ± SD	D	Mean ± SD	D	Min	Max	Min	Max
Males															
10 to <16.5	26	43.5 ± 11.6	L	1.36 ± 0.24	L	1474 ± 287	L	89 ± 36	L	2488 ± 635	L	0.15	0.32	0.28	0.81
16.5 to <25	25	70.5 ± 6.1	N	1.87 ± 0.10	L	1737 ± 156	N	78 ± 41	N	3132 ± 527	N	0.22	0.31	0.35	0.72
25 to <35	46	71.3 ± 6.8	N	1.88 ± 0.12	N	1740 ± 168	L	0 ± 0		3012 ± 467	L	0.21	0.30	0.36	0.79
35 to <45	34	70.3 ± 6.5	N	1.86 ± 0.11	N	1625 ± 148	L	0 ± 0		3008 ± 386	L	0.19	0.30	0.40	0.66
45 to <65	17	72.3 ± 7.9	N	1.88 ± 0.14	N	1681 ± 309	L	0 ± 0		2697 ± 492	L	0.20	0.36	0.34	0.66
65 to ≤96	50	68.9 ± 6.7	L	1.82 ± 0.11	L	1480 ± 187	L	0 ± 0		2286 ± 437	L	0.17	0.30	0.22	0.60
Females															
10 to <16.5	95	45.2 ± 9.1	L	1.39 ± 0.18	N	1278 ± 150	L	82 ± 25	L	2143 ± 457	L	0.15	0.27	0.20	0.73
16.5 to <25	30	60.6 ± 5.6	L	1.68 ± 0.10	N	1385 ± 141	N	17 ± 39	N	2523 ± 294	N	0.15	0.24	0.30	0.61
25 to <35	88	58.7 ± 6.7	L	1.64 ± 0.12	L	1346 ± 154	N	0 ± 0		2387 ± 373	L	0.15	0.26	0.24	0.68
35 to <45	29	58.9 ± 4.8	N	1.64 ± 0.08	N	1320 ± 114	N	0 ± 0		2441 ± 334	L	0.15	0.22	0.31	0.57
45 to <65	51	58.7 ± 4.9	N	1.63 ± 0.09	N	1211 ± 139	L	0 ± 0		2128 ± 338	N	0.14	0.24	0.23	0.57
65 to ≤96	45	57.2 ± 7.3	L	1.60 ± 0.13	L	1217 ± 152	L	0 ± 0		1729 ± 383	L	0.15	0.25	0.16	0.52

N = normal; L = lognormal. *n* = number of individuals; SD = standard deviation.

^aNormal-weight for children aged 10–19 years with body mass index (BMI) corresponding to 85th percentile or below and adults aged 20–96 years with BMIs between 18.5 and 25 kg/m² (IOM 2002).

^bBEE = basal energy expenditure (i.e. basal metabolic rate expressed on a 24-h basis) measured by indirect calorimetry (IOM 2002).

^cECG = stored daily energy cost for growth (Brochu et al. 2006a).

^dTDEE = total daily energy expenditure. TDEEs were based on ²H₂O and H₂¹⁸O disappearance rates from urine monitored by gas-isotope ratio mass spectrometry during 7- to 21-day periods for free-living individuals (IOM 2002).

^eVO₂β and VO₂α = oxygen consumption rates for individuals at rest and during aggregate daytime activities, respectively, used for the selection of data for the calculation of *H* and *VQ* values. D = best fit distribution (i.e. lognormal or normal) according to the Anderson-Darling goodness-of-fit test performed on individual data for each age group.

physical capacity and opportunities for doing a great deal of demanding exercises (Polgar and Weng 1979; Guyton 1991). No published VE and VO₂ were found for children from 1 to <10 years of age for VO₂ demands within the span of VO₂α. Values for VQβ and VQα for the latter age group were assumed to be the same (i.e. 30.8 ± 0.9), considering the small difference found between VQβ (27.7 ± 3.4, *n* = 145) and VQα (29.9 ± 4.2, *n* = 166) values in older children aged 10 to <16.5 years (Table 3).

Accuracy of energetic measurements

The accuracy of *E*, as well as the BEE values based on the gas exchange of VO₂ and VCO₂ monitored by indirect calorimetry and calculated with the use of the Weir equation (Equation 7), has been shown to vary from +1% to +2% compared with values measured by steady-state direct calorimetry in a sealed chamber (Turel and Alexander 1964). Consequently, *H_p* and *H_F* values calculated based on the Equation 8 are affected by an error ranging from −2% to −1%. During DLW measurements, subjects are advised not to change their usual sources of ingested water for the entire duration of the study. Changing water sources during the isotope elimination period has been found to lead to an increase in the mean error of TDEE values by −8.7% in infants and +5.3% in soldiers (Delany et al. 1988; Jones et al. 1988). However, the mean accuracy of TDEE values from DLW method has been validated against other methods, including metabolic chambers as varying from −1.0%

to +3.3% when the sources of tap water were not modified during the entire period (IDECG 1990). This range of errors also affects the accuracy of ECG values (Brochu et al. 2006a). Therefore, the combined effects of simultaneous minimal and maximal mean errors associated with *H_p*, *H_F* (i.e. −2% to −1%), BEE (i.e. +1% to +2%), TDEE, and ECG values (i.e. −1.0% to +3.3%) on the order of magnitude of physiological daily inhalation rates were determined in the present study.

Physiological daily inhalation rates

Tidal volumes, breathing frequency rates, VE and VO₂ values (e.g. Tabachnik et al. 1981; Colrain et al. 1987; Hudgel et al. 1993; Morrell et al. 1995), systolic and diastolic blood pressures, and heart rates have all been shown to be lower in sleeping subjects compared with their awaken counterparts (e.g. Carrington et al. 2005; Zaregarizi et al. 2007). These findings are in accordance with the reduction of BMR values during sleep. Based on heat production measured in sleeping subjects by direct calorimetry, the sleeping metabolic rates (SMR) were calculated to be 0.960 ± 0.023 times the BMR values in normal-weight (*n* = 86) individuals (Benedict and Carpenter 1910; Buskirk et al. 1960; Bessard et al. 1983; Schutz et al. 1984; Shapiro et al. 1984; Ravussin et al. 1985; Garby et al. 1987). This correcting factor (referred to as *F_{sleep}*) affecting BEE values as well as the minimal and maximal *F_{sleep}* values of 0.870 and 1.039 were integrated into the following

Table 3. Ventilatory equivalents for healthy individuals aged 2 h to 96 years at rest and during aggregate daytime activities.

Age groups for both genders (years)	Ventilatory equivalents (L of air inhaled/L of O ₂ consumed) ^a															
	During resting conditions ^b (VQβ)				During the aggregate daytime activities ^b (VQα)				Below the anaerobic threshold				During anaerobiosis			
									VO ₂ ^c				VO ₂ ^c			
	n	Mean ± SD	Min	Max	n	Mean ± SD	Min	Max	n	Mean ± SD	Min	Max	n	Mean ± SD	Min	Max
<1	131	30.2±7.6	16.7	60.8	Same values as those for VQβ				Not applicable				Not applicable			
1 to <10	35	30.8±0.9	25.4	46.6	Same values as those for VQβ				27	26.6±0.9	0.54	0.70	88	36.9±5.2	0.86	2.62
10 to <16.5	145	27.7±3.4	17.1	39.4	166	29.9±4.2	18.9	49.2	23	30.0±0.4	0.76	0.92	1282	37.6±2.1	1.50	4.47
16.5 to <25	114	27.4±4.8	14.4	47.4	85	32.2±6.1	21.0	100.5	459	26.9±2.0	0.72	1.81	818	35.3±2.3	3.00	5.63
25 to <35	133	32.2±3.1	18.0	64.0	318	32.6±4.7	15.7	84.6	390	29.8±2.8	0.80	1.78	535	33.6±1.5	3.01	5.18
35 to <45	60	30.6±2.2	22.1	48.0	47	33.1±8.6	15.3	91.5	205	26.1±1.9	0.75	1.75	125	37.7±1.5	3.14	4.23
45 to ≤96	38	30.6±2.6	22.3	40.8	59	33.7±7.2	16.0	76.5	89	28.7±2.1	0.77	1.17	2736	35.7±1.0	1.51	4.94

n = number of individuals; SD = standard deviation; Min = minimal value; Max = maximal value.

^aVQ = ratio of the minute ventilation rate (VE in L/min at BTPS) to the oxygen uptake (VO₂ in L/min at STPD). The simultaneous VE and VO₂ measurements used for VQ calculations were taken from different studies, which are cited in the appendix.

^bVO₂ values for VQβ and VQα vary from 0.06 to 0.36 and 0.06 to 0.81 L/min, respectively (Tables 1 and 2).

^cOxygen consumption rate (in L/min).

equation in order to determine the SMR values (in kcal/min) for subjects during sleep in the supine position:

$$SMR = \left[\frac{(BEE \times F_{\text{sleep}}) + ECG}{1440} \right] \quad (9)$$

Values for physiological daily inhalation rates (m³/day) were then calculated by using the following expression:

$$PDIR = \left[\frac{(SMR \times H_F \times VQ\beta \times Sld) + (E\alpha \times H_p \times VQ\alpha \times (24 - Sld))}{0.060} \right] \times 0.06 \quad (10)$$

where 0.060 is the combined conversion factor from hours to minutes and liters (L) to cubic meters (m³).

Values for physiological daily inhalation rates expressed per unit of BSA were determined by using the BSA values calculated on the basis of height (cm) and weight (kg) values as follows (Mosteller 1987):

$$BSA = \left[\frac{\text{height} \times \text{weight}}{3600} \right]^{0.5} \quad (11)$$

Sleep durations

Sld from the literature were used in this study regardless of the under-, normal-, overweight, and obese proportions of individuals in the different cohorts. However, several publications suggest that overweight and obese children and adults have shorter night sleep compared with their normal-weight counterparts (Taheri *et al.* 2004; Cizza *et al.* 2005; Gangwisch *et al.* 2005; Vorona *et al.* 2005; Beebe *et al.* 2006; Kohatsu *et al.* 2006; Patel *et al.* 2006; Taheri 2006; Bjorvatn *et al.* 2007; Seicean *et al.* 2007). On the contrary, some publications challenge this view (Koçoglu *et al.* 2003; Gibson *et al.* 2004; Hasler *et al.* 2004; Eisenmann *et al.* 2006). To dispel the influence of this ambiguity on inhalation values, two sets of daily inhalation rates were calculated and compared.

A first set of physiological daily inhalation rates was calculated by using the Sld reported in Bernstein *et al.* (2001) and Eisenmann *et al.* (2006) for small cohorts of subjects composed of known proportions of normal-weight, overweight, and obese individuals (i.e. 10.9% of boys and 13.6% of girls aged 7.5 to 10.9 years, 10.7% of males and 11.9% of females aged 11–16.5 years were overweight or obese; in adults aged 35–74 years, 45% of males and 24% of females were overweight, 9% of males and 13% of females were obese). This set of values was compared with a second set of inhalation rates that was calculated when Sld for percentages of overweight/obese individuals in both cohorts were decreased by 25%. This process of calculation corresponds to the worst case scenario based on data reported in the literature. Sld for 60% of overweight/obese children were decreased by 25% based on published values that indicate that 13.5–57.6% of overweight/obese children aged 7.5–16.5 years ($n=6426$) have sleep deprivation varying from 13.1% to 21.9% (Eisenmann *et al.* 2006; Seicean *et al.* 2007). Sld for 35% of overweight adults and 55% of their obese counterparts were decreased by 25% considering the fact that 27.8–35.1% of overweight adults and 29.3% to 53.1% of their obese counterparts ($n=96,570$) aged 32–86 years (Gangwisch *et al.* 2005; Kohatsu *et al.* 2006; Patel *et al.* 2006; Bjorvatn *et al.* 2007) have Sld 14.3–25.4% and 16.4–26.2%, respectively, shorter than the healthy baseline of 7 h per night (Kripke *et al.* 2002; Patel *et al.* 2004; Cizza *et al.* 2005; Gangwisch *et al.* 2005; Seicean *et al.* 2007).

Statistical analysis

Means, SD values, and distribution percentiles were calculated for all values, which were grouped by age with >30 subjects per group in order to optimize the probability of achieving a normal distribution for each age group, as formally recommended according to the central limit theorem (Feller 1945; Trotter 1959; Rice 1995). Monte

Carlo simulations were used in order to take into account SD values into mean calculations and different physiological equations. Each calculation process was based on random sampling involving 10,000 iterations. Log normal distributions of Sld were used into the calculation process of physiological daily inhalation rates based on data reported in Knutson and Lauderdale (2007) and Seicean *et al.* (2007). Distributions of other parameters were defined to be either lognormal or normal according to the Anderson-Darling goodness-of-fit tests performed on individual data (Tables 1-4). The same statistical test was used to define the best fit distributions of the resulting physiological daily inhalation rates per age group. The number of individual observations in fasting subjects reported in Gibney *et al.* (2003) and Shepherd *et al.* (2007) was insufficient ($n=8$) for the use of the Anderson-Darling test. Therefore, individual H_p ($n=102$) values for subjects in the supine position were statically tested in order to characterize the distribution type for the H_F value during nighttime sleep.

Results

Mean and SD values for body weights, BSA, BEE, ECG, and TDEE as well as lower and upper limits of $VO_2\beta$ and $VO_2\alpha$ are presented in Tables 1 and 2, while those for

Sld are reported in Table 5. Mean H_p values resulting from nutrient intakes in different countries are given in Tables 6-8. Those for mean and SD values and distribution percentiles of $VQ\beta$ and $VQ\alpha$ values as well as mean and SD values for VQ ratios below the anaerobic threshold and during anaerobiosis are presented in Table 3. Means and distribution percentiles of physiological daily inhalation rates in normal-weight males and females aged 2.6 months to 96 years are given in Tables 9 and 10 respectively. Mean values for daily inhalation rates as a function of age are presented in Figures 1-3.

Results of Anderson-Darling goodness-of-fit tests on anthropometric and energetic values are reported in Tables 1 and 2, while those on respiratory parameters are given in Table 4. Finally, physiological daily inhalation rates for all age groups in m^3/day , $m^3/\text{kg-day}$, as well as $m^3/m^2\text{-day}$ better fit with lognormal distributions except for those in m^3/day for girls aged 1 to <2 years, which better fit with a normal distribution (data not shown in tables).

Values for H_p based on dietary intakes were found to vary between 0.203 and 0.208 L of O_2/kcal in 17 countries (Table 6), albeit North-American values range from 0.206 to 0.208 L of O_2/kcal . H_p values for the Canadian population range from 0.205 to 0.207 L of O_2/kcal for in term infants after birth and remain relatively constant

Table 4. Distribution type of parameters used in the calculation process of ventilatory equivalents, oxygen uptake factors, and physiological daily inhalation rates.

Parameter	Acronym ^a	<i>n</i>	Age (years)	Distribution
Oxygen consumption rate (L/min)	$VO_2\beta$	337 ^b	0.22 to 96	Normal
	$VO_2\alpha$	307 ^b	1 to 96	Lognormal
	$VO_{2\text{Sub-anaerobiosis}}$	682 ^b	1 to 96	Lognormal
	$VO_{2\text{Anaerobiosis}}$	296 ^b	1 to 96	Lognormal
Carbon dioxide production (L/min)	$VCO_2\beta$	162 ^b	0.22 to 96	Normal
	$VCO_2\alpha$	117 ^b	1 to 96	Lognormal
	$VE\beta$	131 ^b	0.22 to <1	Normal
Minute ventilation rate (L/min)	$VE\beta$	49 ^b	1 to 96	Lognormal
	$VE\alpha$	141 ^b	1 to 96	Lognormal
	$VE_{\text{Sub-anaerobiosis}}$	682 ^b	1 to 96	Lognormal
	$VE_{\text{Anaerobiosis}}$	296 ^b	1 to 96	Lognormal
	$RER\beta^c$	162 ^b	1 to 96	Lognormal
Respiratory exchange ratio (unitless)	$RER\alpha^c$	117 ^b	1 to 96	Lognormal
	$VQ\beta^d$	280 ^b	0.22 to 96	Lognormal
Ventilatory equivalent ratio (unitless)	$VQ\alpha^d$	141 ^b	1 to 96	Lognormal
	H_F^e	102 ^b	0.22 to 96	Normal
Oxygen uptake factor (L of O_2/kcal)	H_p^f	229 ^b	0.22 to 96	Lognormal
	Sld	2055 ^g	0.22 to 96	Lognormal

n = number of individual data on which the best fit distribution (i.e. lognormal or normal) has been defined.

^a β = for subjects at rest. α = during the aggregate daytime activities of subjects.

^bBest fit distribution (i.e. lognormal or normal) according to the Anderson-Darling goodness-of-fit test performed on individual data taken from studies cited in the appendix and Johnson *et al.* (1960), Reeves *et al.* (1961), Åstrand *et al.* (1964), Frick and Somer (1964), Emirgil *et al.* (1967), Hermansen *et al.* (1970), Jones *et al.* (1970), Pernow and Saltin (1971), and Capderou *et al.* (1997).

^c $RER = VCO_2/VO_2$ ratio.

^d $VQ = VE/VO_2$ ratio.

^eDuring fasting phase.

^fDuring postprandial phase.

^gLognormal distribution based on data reported in Knutson and Lauderdale (2007) and Seicean *et al.* (2007).

Table 5. Sleep duration in healthy individuals aged 2.6 months to 96 years.

Gender and age group (years)	Sleep duration (h/day)	
	<i>n</i>	Mean \pm SD
For both genders		
0.22 to <0.5	456 ^a	14.2 \pm 1.9
0.5 to <1	916 ^a	13.9 \pm 1.0
1 to <2	912 ^a	13.4 \pm 0.8
2 to <5	1361 ^a	11.9 \pm 0.6
5 to <7	900 ^a	10.8 \pm 0.5
Males		
7 to <10	919 ^b	9.9 \pm 1.2
10 to <16.5	2284 ^b	9.2 \pm 0.8
16.5 to <25	552 ^c	8.0 \pm 1.2
25 to <35	127 ^c	8.0 \pm 2.0
35 to <45	670 ^d	7.2 \pm 0.7
45 to <65	1192 ^d	8.0 \pm 0.5
65 to \leq 96	366 ^d	8.8 \pm 0.7
Females		
7 to <10	953 ^b	10.2 \pm 1.0
10 to <16.5	2168 ^b	9.3 \pm 0.8
16.5 to <25	712 ^c	8.5 \pm 1.1
25 to <35	172 ^c	8.4 \pm 1.6
35 to <45	784 ^d	8.1 \pm 0.7
45 to <65	1196 ^d	8.2 \pm 0.5
65 to \leq 96	376 ^d	9.1 \pm 0.7

^aIglowstein *et al.* (2003).^bEisenmann *et al.* (2006).^cAdams (2006).^dAdams (2006) and Bernstein *et al.* (2001).

(variation of values $\leq 0.5\%$) into advanced age (Table 7). Values for H_p were confirmed to almost always be identical between males and females of the same age living in the same country. Variations observed were consistently $<0.4\%$ (Table 8). Values of 0.206, 0.207 and 0.209 L of O_2 /kcal were calculated for the 10th, 50th, and 90th percentiles based on Canadian nutrient intake contributions observed and compiled by Brault-Dubuc and Mongeau (1989) over a 10-year span ($n=747$). H_p values were calculated to be 0.206, 0.207, 0.207 L of O_2 /kcal for underweight ($n=14$), normal-weight ($n=25$), and obese adults ($n=18$), respectively, based on typical German diet (Bosy-Westphal *et al.* 2004). H_p values for black ($n=246$) and white Americans ($n=703$), calculated in this study, based on their nutrient intakes (Morisson *et al.* 1980) vary by $<0.5\%$.

Results of H_p and H_F values calculated based on simultaneous VO_2 and VCO_2 measurements are not shown in tables. Values for an H_F of 0.205 ± 0.003 , 0.206 ± 0.003 , and 0.207 ± 0.003 L of O_2 /kcal for subjects at rest in a semi-recumbent (Müller *et al.* 1989; $n=5$), almost supine (Saltzman and Salzano 1971; $n=20$) and supine position (Gibney *et al.* 2003; $n=6$) were calculated with VO_2 , E , and RER values varying from 0.225 ± 0.035 to 0.307 ± 0.044 L/min, 1.09 ± 0.05 to 1.47 ± 0.07 kcal/min, and 0.802 ± 0.057 to 0.858 ± 0.072 , respectively. A mean H_F value of 0.205 ± 0.001 L of O_2 /kcal was also calculated for adults aged 23–30 years ($n=27$) performing exercise in the

upright position below the anaerobic threshold (De Bock *et al.* 2005; VO_2 of 2.83 ± 0.05 L/min, VCO_2 of 2.37 ± 0.05 L/min, E of 13.41 ± 0.21 kcal/min, RER of 0.838 ± 0.023 with minimal and maximal values of 0.759 and 0.928, respectively). These results show that the level of exertions in fasting subjects (thus VO_2 demands at rest or during exercise below the anaerobic threshold), and their positions (i.e. upright or supine position) during measurements had a negligible effect on their H_F values (by $<1\%$). Consequently, H_F and H_p values for nighttime sleep and the aggregate daytime activities, respectively, were calculated by using VO_2 and VCO_2 values measured in healthy subjects while performing activities with VO_2 demands that were within the entire span of $VO_{2\beta}$ and $VO_{2\alpha}$ values varying from 0.06 to 0.79 L/min (Tables 1 and 2) regardless of their positions during the experimental protocols. Values for H_F of 0.2057 ± 0.0018 L of O_2 /kcal ($n=31$) and H_p of 0.2059 ± 0.0019 L of O_2 /kcal ($n=1245$) were then calculated. The H_p value was calculated by using published data for individuals aged 2 h to 73 years ($n=327$) in the supine position and 8.8–81 years ($n=918$) in the upright position (Tenney and Miller 1956; Baker *et al.* 1957; Spurr *et al.* 1957; Emirgil *et al.* 1967; Pernow and Saltin 1971; Oren *et al.* 1981; Allen *et al.* 1984; Capderou *et al.* 1997; Treuth *et al.* 1998; Gisolf *et al.* 2003; Cade *et al.* 2004; Shiou-Liang *et al.* 2005; other references are underlined in the appendix). During the postprandial phase, VO_2 , E , and RER values of 0.184 ± 0.011 L/min, 0.90 ± 0.04 kcal/min, and 0.866 ± 0.074 , respectively, were calculated for individuals in the supine position, compared with a VO_2 of 0.291 ± 0.013 L/min, an E of 1.41 ± 0.05 kcal/min, and a RER of 0.817 ± 0.050 for subjects in the upright position.

The worst case scenario of decreased Sld in overweight/obese subjects has reduced the global physiological daily inhalation rates of entire cohorts of subjects by only 0.03% to 0.17% (data not shown in tables). Initial Sld of 9.9 ± 1.2 ($n=919$), 9.2 ± 0.8 ($n=2284$), 7.8 ± 0.3 h/day ($n=1707$) in males and 10.2 ± 1.0 ($n=953$), 9.3 ± 0.8 ($n=2168$), 8.2 ± 0.4 h/day ($n=1703$) in females have been published for subjects aged 7.5–10.9, 11–16.5, and 35–74 years, respectively (Bernstein *et al.* 2001; Eisenmann *et al.* 2006). Classified in the same order, initial Sld values of entire cohorts of subjects were decreased to 9.7 ± 1.1 , 9.0 ± 0.8 , 7.3 ± 0.3 h/day for males and 10.0 ± 0.9 , 9.1 ± 0.8 , 7.9 ± 0.3 h/day for females as a result of a 25% reduction in Sld for 60% of overweight/obese children and 35% of overweight as well as 55% of obese adults. Sld specifically for overweight/obese subjects aged 7.5–10.9, 11–16.5, and 35–74 years were decreased to 7.4 ± 0.8 , 6.9 ± 0.6 , 5.8 ± 0.3 h/day in males and 7.6 ± 0.7 , 7.0 ± 0.6 , 6.1 ± 0.3 h/day in females, respectively.

Lower and upper mean errors associated with H_p , H_F (i.e. -2% to -1%), BEE (i.e. $+1\%$ to $+2\%$), TDEE, and ECG values (i.e. -1.0% to $+3.3\%$) affect physiological daily inhalation rates by -2.0 to -1.0 , -0.08 to -0.01 , -1.0 to $+3.4$, and -0.2 to $+0.7\%$, respectively. Simultaneous maximal mean errors associated with H_p , H_F values (-1%), BEE ($+2\%$), ECG, and TDEE values ($+3.3\%$) increase

Table 6. Postprandial oxygen uptake factor resulting from daily nutrient intakes for all ages by country.

Age (years)	n	Nutrient intake contributions (%)			Oxygen uptake factor ^b (L of O ₂ /kcal)	Country
		Protein	Fat	COH ^a		
1 to 9	1442	17.4	20.6	62.0	0.206	Australia ^c
<1 month to 65+	13,211	17.4	20.6	62.0	0.206	Canada ^d
24 to 74	1010	21.9	15.1	63.0	0.206	China ^e
1 to 24	3147	17.7	21.8	60.5	0.206	Finland ^{fg}
3 to 65+	3003	16.7	38.1	45.3	0.208	France ^h
25 to 27	57	14.1	35.0	48.3	0.207	Germany ⁱ
8.9	116	9.2	22.0	68.0	0.204	Ghana ^f
2 to 8	101	18.5	23.2	58.2	0.206	Greece ^j
2 to 6	99	11.9	24.8	63.3	0.205	India ^k
9 and ≤60	1055	17.6	17.9	64.3	0.205	Italy ^l
40 to 50	351	19.4	15.1	65.5	0.205	Japan ^m
2 to 8; 50 to 69	1225	16.5	26.9	56.0	0.206	Sweden ^{ln}
1.3 to 9	684	13.5	35.3	51.3	0.207	The Netherlands ^{fo}
8.8	114	11.7	16.0	72.0	0.204	The Philippines ^f
0.5 to 1.2	2026	15.4	21.8	64.4	0.205	UK ^p
1 week to 75+	74,275	17.4	34.6	47.6	0.208	USA ^{l,q}
All ages	17,763	9.8	11.4	78.9	0.203	Vietnam ^r

^aCOH = Carbohydrate.^bH_p = postprandial oxygen uptake factor.^cHitchcock et al. (1984) and Jenner et al. (1988).^dNC (1977), Leung et al. (1984), and Brault-Dubuc and Mongeau (1989).^eWoo et al. (1998).^fKnuiman et al. (1983).^gRäsänen et al. (1985), Räsänen et al. (1991), and Räsänen and Ylönen (1992).^hRazanamahefa et al. (2005).ⁱBosy-Westphal et al. (2004).^jNeiderud et al. (1992).^kNarasinga et al. (1982).^lFreudenheim et al. (1993).^mTokudome et al. (1998).ⁿRiboli et al. 1997.^oHoffmans et al. (1986).^pBransby and Fothergill (1954), Margarey and Boulton (1984), Nelson et al. (1990), Payne and Belton (1992), and Ruxton et al. (1996).^qMorisson et al. (1980), Butte and Calloway (1981), Reichman et al. (1981), DHHS (1983), Gross (1983), Butte et al. (1984), USDA (1984), Pao et al. (1985), Oliveria et al. (1992), Simons-Morton et al. (1997), and Bollella et al. (1999).^rThang and Popkin (2004).

daily inhalation values by +2.3%. The inverse scenario is observed with simultaneous minimal mean errors for H_p, H_F (-2%), BEE (+1%), ECG, and TDEE (-1.0%) values affecting physiological daily inhalation rates by -3.0%. The use of SMR instead of BEE values (in Equation 10) has reduced daily inhalation values by only 0.6% to 1.8%. The use of the lowest H value of 0.203 L of O₂/kcal for Vietnamese (n = 17,763) and the highest value of 0.208 L of O₂/kcal for American (n = 74,275) during the postprandial phase (Table 6) could have affected the physiological daily inhalation rates by only -1.2% to -0.7% and +0.5% to +0.9%, respectively, compared with the inhalation values calculated in this study based on H_p value of 0.2059 L of O₂/kcal.

Discussion

All mean and almost all (98%) percentile values of physiological daily inhalation rates calculated in the present article (in m³/day, m³/kg-day, and m³/m²-day) are

higher in males than in females, and are in accordance with Brochu *et al.* (2006a-c). As found in our previous studies, mean daily inhalation values expressed in m³/kg-day follow a logarithmic pattern (Figure 2). Values drop rapidly with increasing age, from 16.5 to <25 years in females (R² = 0.94) and males (R² = 0.96). Then mean physiological daily inhalation rates continue to decrease slowly as age increases up to 65 to 96 years. Mean daily inhalation values in males (0.225 ± 0.059 m³/kg-day) and females (0.202 ± 0.059 m³/kg-day) aged 65-96 years are found to be 61% and 64% lower, respectively, than those for boys (0.572 ± 0.191 m³/kg-day) and girls (0.563 ± 0.180 m³/kg-day) 2.6 to <6 months old. When females and males age from 2.6 months to <16.5 years, body weights increase proportionally more (by 9.2- and 10.6-folds, respectively) than height does (by 2.7- and 2.8-folds, respectively). This results in a moderate increase of BSA values by 5.0- and 5.5-folds, respectively. Beyond these ages, very few changes appear for weight, height, and BSA values. This explains why the

Table 7. Postprandial oxygen uptake factor resulting from daily nutrient intakes for both sexes as a function of age.

Age	n	Nutrient intake contributions (%)			Oxygen uptake factor ^b (L of O ₂ /kcal)
		Protein	Fat	COH ^a	
Breast milk					
1 week ^c	60	18.7	26.3	55.1	0.207
2 weeks ^c	60	15.3	27.9	56.8	0.206
3 weeks ^c	60	13.0	28.4	58.5	0.206
4 weeks ^c	60	11.6	30.6	57.8	0.206
5 weeks ^c	60	11.0	30.4	58.6	0.206
1 month ^d	37	9.0	33.3	57.7	0.206
1 month ^e	10	12.2	34.8	53.0	0.206
6 weeks ^c	60	10.9	31.1	58.0	0.206
7 weeks ^c	60	10.4	30.3	59.3	0.205
8 weeks ^c	60	10.2	29.4	60.4	0.205
2 months ^d	40	8.2	31.7	60.1	0.205
9 to 10 weeks ^c	60	9.9	30.1	60.0	0.205
10 to 12 weeks ^c	60	9.7	29.0	61.3	0.205
3 months ^d	37	7.9	30.4	61.8	0.205
4 months ^d	41	7.5	31.6	60.9	0.205
Formula-fed ^f					
1 to 12 weeks	60	15.3	26.4	58.3	0.206
Liquid and solid food ^g					
<1 month	6	15.7	21.7	62.7	0.205
1 to 2 months	35	15.6	20.6	63.8	0.205
3 to 5 months	65	21.1	15.7	63.3	0.206
6 to 8 month	74	21.1	17.1	61.8	0.206
9 to 11 months	70	19.8	14.5	65.6	0.205
1 to 4 years	1031	18.5	20.9	60.6	0.206
5 to 11 years	1995	16.6	20.7	62.6	0.206
19 to 19 years	2232	17.3	22.8	59.9	0.206
20 to 39 years	2346	18.9	24.0	57.1	0.206
40 to 64 years	2722	18.8	23.2	58.0	0.206
65+ years	1699	18.0	21.7	60.3	0.206

^aCOH = Carbohydrate.^bH_p = postprandial oxygen uptake factor.^cGross (1983).^dButte et al. (1984).^eButte and Calloway (1981).^fGross (1983).^gValues for Canadian individuals (NC 1977).

mean physiological daily inhalation rates expressed in m³/m²-day begin to decrease linearly only as age increases from the age groups of 10 to <16.5 years for males ($R^2=0.92$) and 16.5 to <25 years for females ($R^2=0.94$) up to the age group of 65–96 years (Figure 3). Mean daily inhalation rates for boys 0.22 to <16.5 years old (10.99 ± 3.50 to 11.82 ± 3.50 m³/m²-day) and girls 0.22 to <10 years of age (10.81 ± 3.29 to 10.83 ± 1.84 m³/m²-day) are higher than those for older males and females (8.42 ± 2.13 to 10.93 ± 2.80 and 7.20 ± 1.99 to 9.90 ± 2.50 m³/m² day, respectively). Furthermore, in agreement with our previous study, mean physiological daily inhalation rates in females as well as males aged 25 to <65 years (14.46 ± 3.37 to 20.12 ± 5.03 m³/day and 0.247 ± 0.061 to 0.289 ± 0.077 m³/kg-day) are lower than those for normal-weight pregnant and lactating females

aged 23–55 years, whose values vary from 19.00 ± 9.98 to 22.31 ± 2.50 m³/day and 0.297 ± 0.056 to 0.330 ± 0.069 m³/kg-day (Brochu *et al.* 2006b). Moreover, mean daily inhalation rates in boys (0.428 ± 0.098 to 0.572 ± 0.191 m³/kg-day) and girls (0.395 ± 0.076 to 0.563 ± 0.180 m³/kg-day) aged 0.22 to <10 years are higher than the highest means for under-, normal-, overweight, and obese gravid and breastfeeding females aged 11–55 years of 0.385 ± 0.110 and 0.383 ± 0.064 m³/kg-day, respectively, as reported in Brochu *et al.* (2006b). This is the case in spite of (1) higher VQ means (34.2–36.8) used in the calculation of inhalation rates in pregnant and lactating females compared with those (30.2 and 30.8) in non-gestational and non-lactating individuals and (2) similar mean *H* values (0.21 and 0.206 L/kcal, respectively).

Based on means and percentiles of physiological daily inhalation rates calculated in the present study, children are generally expected to inhale more air pollutants per unit of weight and BSA (i.e. in µg/kg-day and µg/m² day, respectively) than adults during identical exposure concentrations and conditions. The same applies when males are compared with females. The new methodology developed in this study therefore illustrates that some individuals inhale more air on a daily basis (thus more air pollutants) than estimated before. In males 16.5 to <25 years of age, 95th, 97.5th, and 99th percentile values of 28.05, 30.02, and 31.89 m³/day, respectively, were determined. In males 35 to <45 years old, corresponding percentiles were 29.32, 31.84, and 35.40 m³/day, respectively. Values from the 95th to 99th percentile in children younger than 1 year of age vary from 0.806 to 1.105 m³/kg-day in girls and 0.842 to 1.138 m³/kg-day in boys. These percentiles are 2.8- to 4-folds higher than the inhalation estimate of 0.286 m³/kg-day (i.e. 20 m³/day for a 70-kg adult) adopted by the Federal Register Notices (1980). The same nearly applies to the span of values from the 5th to 99th percentiles (0.328–1.138 m³/kg-day) for children aged 0.22 to <7 years, and the 10th to 99th percentiles (0.303–0.712 m³/kg-day) for those from 7 to <10 years old.

The magnitude of human variability in inhalation values, as reflected by the lowest 1st percentile of 0.105 m³/kg-day (data not shown in tables) and the highest 99th percentile of 1.138 m³/kg-day in males and females aged 2.6 months to 96 years (Tables 9 and 10) corresponds to a factor of 10.9. The inter-individual variability factor of 4.8 was also calculated as the ratio of the highest 95th percentile of 0.937 m³/kg-day to the lowest 50th percentile of 0.194 m³/kg-day. Values for lowest percentiles were always observed in elderly females aged 65 to <96 years and the highest percentile was found in boys aged 2.6 to <6 months. Such inter-individual variability factors for inhalation values (i.e. 4.8–10.9) should be evaluated along with the variability in other pharmacokinetic determinants, in order to assess the adequacy of the default uncertainty factor or the human kinetic adjustment factor (HKAf) currently used in health risk assessment (Renwick 2000; WHO 2005).

Table 8. Oxygen uptake factor resulting from nutrient intakes for both sexes in different countries.

Age (years)	Nutrient intake contributions (%)								Oxygen uptake factor ^c (L of O ₂ /kcal)		Country
	Males				Females				Males	Females	
	<i>n</i>	Prot ^a	Fat	COH ^b	<i>n</i>	Prot ^a	Fat	COH ^b			
1	62	19.6	21.6	58.9	63	20.7	23.0	56.3	0.206	0.207	Australia ^d
1.5	72	18.3	21.4	60.3	70	18.8	21.8	59.4	0.206	0.206	Australia ^d
2	74	17.8	22.1	60.1	72	17.4	21.3	61.2	0.206	0.206	Australia ^d
1 to 2	23	18.1	17.4	64.5	23	17.8	15.1	54.8	0.205	0.206	Finland ^e
2	31	14.7	18.7	66.7	31	15.1	18.7	66.2	0.205	0.205	UK ^f
3	73	16.7	20.6	62.7	72	16.8	21.5	61.7	0.206	0.206	Australia ^d
3	31	15.1	18.3	66.5	42	14.8	19.0	66.2	0.205	0.205	UK ^f
3	153	18.2	19.7	62.1	128	18.3	19.7	62.0	0.206	0.206	Finland ^e
4 to 5	128	16.4	19.0	64.6	139	16.7	19.4	63.8	0.205	0.205	UK ^{f,g}
6	139	17.5	19.1	63.4	145	17.7	19.8	62.6	0.205	0.206	Finland ^e
6 to 9	130	17.0	20.0	63.0	116	17.0	21.3	61.7	0.205	0.206	USA ^h
7 to 10	25	13.6	19.5	66.8	26	14.2	18.7	67.1	0.205	0.205	UK ⁱ
9	281	17.6	21.0	61.3	263	17.7	20.9	61.3	0.206	0.206	Finland ^{j,k}
9.0	434	17.0	19.8	63.3	450	17.0	20.0	63.1	0.205	0.205	Australia ^l
9	133	13.8	37.0	50.0	n.d.	n.d.	n.d.	n.d.	0.207	n.d.	Finland ^m
9	117	13.5	38.0	49.0	n.d.	n.d.	n.d.	n.d.	0.207	n.d.	The Netherlands ^m
9	109	13.4	28.0	57.0	n.d.	n.d.	n.d.	n.d.	0.206	n.d.	Italy ^m
9	114	11.7	16.0	72.0	n.d.	n.d.	n.d.	n.d.	0.204	n.d.	The Philippines ^m
9	116	9.2	22.0	68.0	n.d.	n.d.	n.d.	n.d.	0.204	n.d.	Ghana ^m
9 to 11	196	19.4	21.9	58.7	222	18.9	21.4	59.7	0.206	0.206	USA ⁿ
11 to 12	76	16.0	20.3	63.8	67	15.7	21.0	63.3	0.205	0.205	UK ⁱ
12	274	18.0	21.7	60.2	285	17.6	20.8	61.6	0.206	0.206	Finland ^{j,k}
10 to 12	132	18.9	22.1	59.0	147	16.8	20.8	62.4	0.206	0.206	USA ^h
12 to 14	296	19.4	22.6	57.9	295	19.5	22.6	58.0	0.206	0.206	USA ⁿ
15	257	18.4	22.2	59.4	264	17.5	20.7	61.8	0.206	0.206	Finland ^{j,k}
13 to 15	134	18.0	23.0	59.0	110	17.8	22.0	60.2	0.206	0.206	USA ^h
15 to 18	365	20.3	23.4	56.3	374	20.1	23.1	56.8	0.207	0.207	USA ⁿ
18	217	18.3	22.7	59.0	264	17.5	21.1	61.3	0.206	0.206	Finland ^{j,k}
16 to 19	96	19.1	24.1	56.8	84	17.1	22.7	60.2	0.206	0.206	USA ^h
21.0	73	19.1	23.7	57.2	82	17.5	21.5	61.0	0.206	0.206	Finland ^{j,k}
19 to 22	256	22.0	24.6	53.4	300	21.5	24.1	54.4	0.207	0.207	USA ⁿ
23 to 35	791	21.7	24.8	53.5	952	21.6	23.9	54.5	0.207	0.207	USA ^{n,o}
24	59	19.9	24.6	55.4	84	18.2	21.5	60.3	0.207	0.206	Finland ^{j,k}
24 to<35	117	22.2	15.8	62.1	121	23.3	16.5	60.2	0.206	0.206	China ^p
35 to 40	714	22.6	25.6	51.9	838	22.4	24.8	52.9	0.207	0.207	USA ⁿ
35 to 44	129	21.9	15.7	62.4	134	22.3	15.8	61.9	0.206	0.206	China ^p
40 to 50	171	19.6	14.5	65.9	180	19.2	15.7	65.1	0.205	0.205	Japan ^q
45 to 54	124	22.2	15.2	62.6	127	22.3	14.6	63.1	0.206	0.206	China ^p
51 to 64	579	22.8	25.4	51.7	715	22.6	24.4	53.0	0.207	0.207	USA ⁿ
55 to 74	130	20.5	13.8	65.7	128	20.7	13.8	65.5	0.205	0.205	China ^p
≤60	449	18.1	16.8	65.2	497	18.1	16.8	65.1	0.205	0.205	Italy ^r
≤60	1583	20.9	22.0	57.1	1935	20.1	20.3	59.6	0.207	0.206	USA ^{n,r}

n.d. = not determined. ^aProt = Protein. ^bCOH = Carbohydrate. ^cH_p = postprandial oxygen uptake factor. ^dHitchcock et al. (1984). ^eRäsänen and Ylönen (1992). ^fPayne and Belton (1992). ^gMargarey and Boulton (1984). ^hMorisson et al. (1980). ⁱNelson et al. (1990). ^jRäsänen et al. (1985). ^kRäsänen et al. (1991). ^lJenner et al. (1988). ^mKnuiman et al. (1983). ⁿPao et al. (1985). ^oOliveria et al. (1992). ^pWoo et al. (1998). ^qTokudome et al. (1999). ^rFreudenheim et al. (1993).

The use of H_p, H_p VQβ, and VQα values as calculated in the present study does not invalidate the conclusions of our previous studies based on calculations using a VQ of 27 and H of 0.21 L of O₂/kcal as constant values: (1) the aggregate errors (under- and overestimations) of daily inhalation estimates and percentiles (in m³/day and

m³/kg-day) based on published approaches do remain the same (Brochu *et al.* 2006c) and (2) intakes of inhaled air pollutants per unit of body weight (in µg/kg-day) again are expected to be higher in normal-weight males and females compared with their overweight and obese counterparts (Brochu *et al.* 2006a, b).

Table 9. Distribution percentiles of physiological daily inhalation rates for normal-weight males aged 2.6 months to 96 years.

Age group (years)	Physiological daily inhalation rates ^a										
	Mean \pm SD	Percentiles									
(m ³ /day)		2.5th	5th	10th	25th	50th	75th	90th	95th	97.5th	99th
0.22 to <0.5	3.76 \pm 1.15	2.02	2.20	2.44	2.92	3.57	4.40	5.31	5.97	6.47	7.26
0.5 to <1	4.66 \pm 1.34	2.61	2.82	3.11	3.68	4.46	5.45	6.47	7.13	7.74	8.48
1 to <2	5.68 \pm 0.85	4.24	4.39	4.61	5.05	5.61	6.25	6.86	7.20	7.45	7.75
2 to <5	7.35 \pm 1.39	5.04	5.25	5.57	6.27	7.23	8.31	9.33	9.87	10.24	10.54
5 to <7	9.04 \pm 1.21	6.95	7.21	7.54	8.17	8.94	9.81	10.64	11.18	11.68	12.28
7 to <10	11.17 \pm 1.89	8.14	8.42	8.84	9.74	10.96	12.35	13.82	14.69	15.40	16.21
10 to <16.5	15.64 \pm 3.87	9.82	10.40	11.16	12.74	15.06	17.92	21.04	22.84	24.54	26.72
16.5 to <25	20.39 \pm 4.26	13.30	14.15	15.22	17.37	20.04	22.96	25.93	28.05	30.02	31.89
25 to <35	20.00 \pm 3.78	13.84	14.54	15.52	17.30	19.55	22.27	25.15	27.00	28.52	30.54
35 to <45	20.12 \pm 5.03	12.39	13.24	14.33	16.50	19.41	22.97	26.71	29.32	31.84	35.40
45 to <65	18.41 \pm 4.25	11.86	12.60	13.51	15.30	17.80	20.90	24.05	26.39	28.33	30.75
65 to \leq 96	15.25 \pm 3.78	9.44	10.06	10.90	12.47	14.73	17.50	20.27	22.12	23.91	26.05
(m ³ /kg day) ^b											
0.22 to <0.5	0.572 \pm 0.191	0.290	0.317	0.356	0.433	0.541	0.677	0.828	0.937	1.040	1.138
0.5 to <1	0.536 \pm 0.166	0.288	0.312	0.344	0.414	0.509	0.634	0.759	0.842	0.922	1.015
1 to <2	0.537 \pm 0.095	0.379	0.397	0.420	0.467	0.527	0.599	0.666	0.708	0.747	0.787
2 to <5	0.493 \pm 0.125	0.297	0.317	0.345	0.400	0.477	0.568	0.663	0.726	0.777	0.845
5 to <7	0.463 \pm 0.077	0.332	0.349	0.368	0.407	0.456	0.511	0.564	0.597	0.631	0.668
7 to <10	0.428 \pm 0.098	0.275	0.290	0.312	0.357	0.416	0.485	0.560	0.609	0.653	0.712
10 to <16.5	0.383 \pm 0.131	0.191	0.211	0.237	0.288	0.362	0.454	0.556	0.628	0.702	0.790
16.5 to <25	0.290 \pm 0.065	0.184	0.197	0.213	0.244	0.283	0.330	0.377	0.406	0.435	0.473
25 to <35	0.282 \pm 0.059	0.187	0.198	0.212	0.239	0.275	0.317	0.361	0.390	0.417	0.445
35 to <45	0.289 \pm 0.077	0.173	0.185	0.203	0.234	0.278	0.333	0.389	0.429	0.470	0.523
45 to <65	0.259 \pm 0.065	0.161	0.171	0.184	0.212	0.249	0.296	0.346	0.378	0.408	0.449
65 to \leq 96	0.225 \pm 0.059	0.134	0.144	0.157	0.182	0.216	0.259	0.303	0.333	0.360	0.400
(m ³ /m ² day) ^b											
0.22 to <0.5	10.99 \pm 3.50	5.74	6.26	7.00	8.46	10.44	12.97	15.76	17.60	19.49	21.51
0.5 to <1	11.24 \pm 3.34	6.15	6.69	7.41	8.80	10.74	13.15	15.70	17.42	18.96	21.15
1 to <2	11.68 \pm 1.91	8.42	8.83	9.30	10.28	11.51	12.91	14.25	15.06	15.79	16.59
2 to <5	11.54 \pm 2.61	7.32	7.78	8.40	9.58	11.26	13.18	15.11	16.27	17.36	18.52
5 to <7	11.53 \pm 1.72	8.59	8.96	9.43	10.30	11.39	12.61	13.83	14.58	15.29	16.11
7 to <10	11.55 \pm 2.27	7.94	8.33	8.86	9.88	11.28	12.94	14.61	15.74	16.71	17.75
10 to <16.5	11.82 \pm 3.50	6.64	7.13	7.83	9.26	11.22	13.77	16.61	18.44	20.17	22.29
16.5 to <25	10.92 \pm 2.35	7.02	7.48	8.08	9.23	10.73	12.33	13.96	15.11	16.22	17.45
25 to <35	10.64 \pm 2.12	7.21	7.63	8.12	9.12	10.40	11.89	13.52	14.50	15.43	16.50
35 to <45	10.93 \pm 2.80	6.63	7.11	7.73	8.90	10.53	12.48	14.64	16.05	17.43	19.31
45 to <65	9.88 \pm 2.36	6.28	6.63	7.15	8.16	9.56	11.25	12.99	14.24	15.32	16.92
65 to \leq 96	8.42 \pm 2.13	5.14	5.50	5.95	6.86	8.12	9.67	11.23	12.25	13.29	14.78

SD = standard deviation.

^aDaily inhalation rates = $[(\text{SMR} \times H_F \times \text{VQ}\beta \times \text{Sld}) + (\text{Ea} \times H_p \times \text{VQ}\alpha) \times (24 - \text{Sld})] \times 0.06$, and $\text{SMR} = [(\text{BEE} \times F_{\text{sleep}}) + \text{ECG}] / 1440$, $\text{Ea} = [(\text{TDEE} - \text{BEE}) / ((24 - \text{Sld}) \times 60)] + (\text{BEE} + \text{ECG}) / 1440$. BEE, ECG, TDEE (kcal/day) and Sld (h/day) are defined and given in Tables 1, 2 and 5. VQ β and VQ α (unitless) are defined and reported in Table 3. SMR = sleeping metabolic rate (kcal/min). F_{sleep} is a correcting factor of BEE values. $F_{\text{sleep}} = 0.960 \pm 0.023$, minimum = 0.870, maximum = 1.039. H_F and H_p = oxygen uptake factor during fasting and postprandial phases, respectively (L of O₂/kcal). $H_F = 0.2057 \pm 0.0018$ L/kcal, minimum = 0.198 L/kcal, maximum = 0.214 L/kcal. $H_p = 0.2059 \pm 0.0019$ L/kcal, minimum of 0.199 L/kcal, maximum of 0.221 L/kcal.

^bDaily inhalation rates were divided by body weights and body surface areas reported in Tables 1 and 2 in order to obtain values expressed in m³/kg min and m³/m² min, respectively.

H values

High intakes of carbohydrates and a low level of proteins ingested led to lower H_p values (0.203–0.204 L of O₂/kcal) in subjects ($n = 17\,993$) living in Ghana, the Philippines, and Vietnam, compared with those from other countries (0.205–0.208 L of O₂/kcal; $n = 101,686$). However, the

magnitude of H_p values is unaffected by an individuals' age, gender, or BMI for subjects living in a given country ($n = 119\,679$). Rather, it is the variability of the food intake components that determines the magnitude of H_p values. However, such variability is found to have little effect on the magnitude of physiological daily inhalation

Table 10. Distribution percentiles of physiological daily inhalation rates for normal-weight females aged 2.6 months to 96 years.

Age group (years)	Physiological daily inhalation rates ^a										
	Mean ± SD	Percentiles									
		2.5th	5th	10th	25th	50th	75th	90th	95th	97.5th	99th
(m ³ /day)											
0.22 to <0.5	3.63±1.07	2.03	2.19	2.41	2.86	3.47	4.25	5.07	5.64	6.15	6.72
0.5 to <1	4.30±1.26	2.35	2.57	2.84	3.37	4.13	5.02	6.00	6.62	7.24	8.07
1 to <2	5.43±0.90	3.79	3.97	4.22	4.76	5.41	6.07	6.64	6.96	7.18	7.39
2 to <5	6.90±1.25	4.87	5.06	5.34	5.94	6.77	7.72	8.62	9.20	9.62	9.99
5 to <7	8.59±1.12	6.66	6.88	7.19	7.75	8.50	9.32	10.10	10.58	11.03	11.49
7 to <10	10.71±1.62	7.94	8.27	8.71	9.54	10.57	11.75	12.89	13.64	14.28	14.94
10 to <16.5	13.32±3.06	8.49	9.02	9.67	11.02	12.98	15.17	17.43	18.96	20.35	21.92
16.5 to <25	16.46±3.21	11.19	11.76	12.61	14.16	16.13	18.38	20.69	22.20	23.59	25.22
25 to <35	15.82±3.05	10.83	11.38	12.11	13.65	15.52	17.64	19.88	21.34	22.73	24.35
35 to <45	16.21±4.02	9.99	10.69	11.55	13.31	15.61	18.51	21.68	23.55	25.57	27.90
45 to <65	14.46±3.37	9.04	9.67	10.45	12.05	14.08	16.47	18.91	20.49	22.10	24.00
65 to ≤96	11.51±3.04	6.84	7.32	7.97	9.30	11.07	13.25	15.56	17.29	18.62	20.54
(m ³ /kg day) ^b											
0.22 to <0.5	0.563±0.180	0.299	0.326	0.360	0.431	0.534	0.662	0.807	0.897	0.994	1.105
0.5 to <1	0.510±0.159	0.269	0.295	0.329	0.393	0.486	0.601	0.722	0.806	0.882	0.979
1 to <2	0.516±0.105	0.336	0.356	0.384	0.438	0.510	0.582	0.659	0.704	0.740	0.785
2 to <5	0.492±0.124	0.288	0.311	0.341	0.400	0.480	0.568	0.661	0.716	0.766	0.826
5 to <7	0.441±0.076	0.313	0.328	0.349	0.386	0.434	0.488	0.545	0.579	0.609	0.642
7 to <10	0.395±0.076	0.267	0.284	0.303	0.340	0.388	0.443	0.497	0.531	0.564	0.601
10 to <16.5	0.306±0.089	0.170	0.185	0.204	0.241	0.293	0.358	0.427	0.471	0.514	0.566
16.5 to <25	0.275±0.059	0.180	0.190	0.206	0.234	0.269	0.310	0.352	0.380	0.408	0.444
25 to <35	0.273±0.060	0.176	0.187	0.201	0.230	0.266	0.310	0.354	0.383	0.410	0.443
35 to <45	0.277±0.072	0.166	0.179	0.194	0.225	0.266	0.318	0.373	0.410	0.443	0.480
45 to <65	0.247±0.061	0.150	0.161	0.176	0.203	0.239	0.282	0.328	0.358	0.387	0.420
65 to ≤96	0.202±0.059	0.114	0.124	0.136	0.160	0.194	0.235	0.281	0.311	0.344	0.385
(m ³ /m ² day) ^b											
0.22 to <0.5	10.81±3.29	5.90	6.38	7.02	8.42	10.29	12.62	15.29	17.03	18.65	20.37
0.5 to <1	10.55±3.18	5.71	6.22	6.85	8.23	10.08	12.40	14.77	16.40	17.99	20.00
1 to <2	11.14±2.06	7.47	7.91	8.49	9.61	11.04	12.55	13.93	14.74	15.40	16.10
2 to <5	11.24±2.53	7.12	7.59	8.19	9.36	10.95	12.84	14.66	15.89	16.97	18.04
5 to <7	10.98±1.67	8.12	8.49	8.93	9.77	10.84	12.03	13.24	14.00	14.60	15.41
7 to <10	10.83±1.84	7.68	8.10	8.56	9.49	10.67	12.00	13.29	14.13	14.81	15.58
10 to <16.5	9.67±2.50	5.84	6.23	6.77	7.83	9.35	11.14	13.03	14.29	15.48	16.88
16.5 to <25	9.84±2.00	6.59	6.97	7.45	8.41	9.61	11.00	12.51	13.43	14.35	15.35
25 to <35	9.73±2.01	6.43	6.84	7.33	8.30	9.52	10.93	12.41	13.32	14.27	15.27
35 to <45	9.90±2.50	6.05	6.48	7.01	8.11	9.56	11.32	13.26	14.53	15.76	17.15
45 to <65	8.88±2.12	5.46	5.87	6.35	7.35	8.62	10.12	11.69	12.75	13.67	15.00
65 to ≤96	7.20±1.99	4.22	4.52	4.92	5.76	6.92	8.33	9.87	10.88	11.93	13.15


SD = standard deviation.

^aDaily inhalation rates = [(SMR × H_F × VQβ × Sld) + (Eα × H_p × VQα) × (24 – Sld)] × 0.06, and SMR = [(BEE × F_{sleep}) + ECG]/1440, Eα = [(TDEE – BEE)/((24 – Sld) × 60)] + (BEE + ECG)/1440. BEE, ECG, TDEE (kcal/day) and Sld (h/day) are defined and given in Tables 1, 2 and 5. VQβ and VQα (unitless) are defined and reported in Table 3. SMR = sleeping metabolic rate (kcal/min). F_{sleep} is a correcting factor of BEE values. F_{sleep} = 0.960 ± 0.023, minimum = 0.870, maximum = 1.039. H_F and H_p = oxygen uptake factor during fasting and postprandial phases, respectively (L of O₂/kcal). H_F = 0.2057 ± 0.0018 L/kcal, minimum of 0.198 L/kcal, maximum of 0.214 L/kcal. H_p = 0.2059 ± 0.0019 L/kcal, minimum of 0.199 L/kcal, maximum of 0.221 L/kcal.

^bDaily inhalation rates were divided by body weights and body surface areas reported in Tables 1 and 2 in order to obtain values expressed in m³/kg min and m³/m² min, respectively.


rates. The use of the lowest H_p value of 0.203 L of O₂/kcal for Vietnamese subjects (n = 17,763) and the highest H_p value of 0.208 L of O₂/kcal for American and French subjects (n = 77,278), instead of the H_p value of 0.2059 L of O₂/kcal that was used in this study, would have changed the physiological daily inhalation rates by only –1.2% to

+0.9%. This is due to the fact that H_p and H_F values (i.e. 0.2059 and 0.2057 L of O₂/kcal, respectively) both rest in the middle of the span between the lower Vietnamese (i.e. 0.203 L of O₂/kcal) and higher American (i.e. 0.208 L of O₂/kcal) values. Several thousand sets of VO₂ and VCO₂ values (data not shown in tables; n = 6696) measured in



Plotted values are for midpoint ages of the age cohorts reported in Tables 9 and 10.
Males = solid line; Females = dotted line.

Figure 1. Mean daily inhalation rates (m^3/day) in normal-weight males and females as a function of age.




Plotted values are for midpoint ages of the age cohorts reported in Tables 9 and 10.
Males = solid line; Females = dotted line.

Figure 2. Mean daily inhalation rates ($\text{m}^3/\text{kg day}$) in normal-weight males and females as a function of age.

subjects during strenuous exercise when consuming higher oxygen rates ($0.82\text{--}5.48\text{ L/min}$) than upper $\text{VO}_{2\beta}$ and $\text{VO}_{2\alpha}$ limits would have biased H values that were included in the present study.

VQ values

For a given age group, VQ values during anaerobiosis were found to be higher than values for $\text{VQ}\beta$, $\text{VQ}\alpha$, and VQ for VO_2 demands below the anaerobic threshold ranging from 0.54 to 1.81 L/min . Former VQ values were calculated by using VE values measured in subjects performing strenuous exercises during high oxygen uptake rates varying from 0.86 to 4.47 L/min in children aged 1 to <16.5 years and 3.00 to 5.63 L/min in individuals $16.5\text{--}96$ years of age, respectively. During such periods of exertion, the aerobic metabolism becomes inadequate to supply all energy required and is compensated by the anaerobic metabolism (Guyton 1991). However, these punctual VE values as well as those used for the calculation of VQ values below the anaerobic threshold have little influence on physiological daily inhalation rates, since $\text{VO}_{2\alpha}$ values during the aggregate daytime activities for subjects



Plotted values are for midpoint ages of the age cohorts reported in Tables 9 and 10.
Males = solid line; Females = dotted line.

Figure 3. Mean daily inhalation rates ($\text{m}^3/\text{m}^2 \text{ day}$) in normal-weight males and females as a function of age.

aged 2.6 months to 96 years were found to vary only from 0.06 to 0.81 L/min . The performance of activities under anaerobic conditions can be considered to correspond, in the reality of each day, to sufficiently rare events of short durations; the latter are therefore diluted in the large aerobic process of oxygenation, which is continuously effective during the aggregate daytime activities as well as on a 24-h basis. Consequently, values for VQ during anaerobiosis would have overestimated physiological daily inhalation rates, while most of those during sub-anaerobiosis would have underestimated such rates.

Conclusion

This study presents an exhaustive compilation and critical analysis of a wide range of published data related to H and VQ values. It supports the establishments of solid bases for the appropriate selection and use of input data in the determination of daily inhalation rates. By the same occasion, it contributes to improve our previous procedure based on DLW measurements (Brochu *et al.* 2006a–c) due to the fact that it is now possible to determine and integrate nighttime and daytime respiratory parameters into the physiological daily inhalation calculation process. Only data measured in healthy subjects during VO_2 demands within the span of $\text{VO}_{2\beta}$ and $\text{VO}_{2\alpha}$ values based on DLW measurements were used in the present study in order to determine H_F and $\text{VQ}\beta$ values for nighttime sleep (fasting phase) as well as H_p and $\text{VQ}\alpha$ values for aggregate daytime activities (postprandial phase), respectively. This innovative strategy has allowed for the exclusion of inadequate published data in the calculation of physiological daily inhalation rates measured in $>19,000$ subjects. Values for H_p , $\text{VQ}\beta$, H_F , and $\text{VQ}\alpha$ were combined into the daily inhalation rates calculation process with BEE from indirect calorimetry measurements ($n = 1235$) as well as ECG and TDEE values based on DLW methodology covering an aggregate period of $>19,000$ days. In the worst case scenario, simultaneous minimal and maximal mean errors associated

with H , BEE, ECG, and TDEE values could have a combined effect varying from -3.0% to $+2.3\%$ on the accuracy of physiological daily inhalation values. This span of potential errors is insignificant compared with those based on time-activity ventilation, food-energy intakes, metabolic equivalents, and Parameter A approaches (Brochu *et al.* 2006c), which vary from -49% to $+122\%$ for some 24-h breathing estimates. Body weight and height, as well as BEE and TDEE values that have been systematically measured for each subject during DLW measurements have assured a precise calculation of inhalation rates per unit of weight and BSA in the present study. Mean and percentile physiological daily inhalation rates expressed in $\text{m}^3/\text{m}^2\text{-day}$ have never been determined before for individuals as a function of age. The information presented strongly suggests that the mean and percentile physiological daily inhalation values reported in this study correspond to the most precise inhalation values (in m^3/day , $\text{m}^3/\text{kg-day}$, and $\text{m}^3/\text{m}^2\text{-day}$) in current literature, and are thereby relevant for use in health risk assessment.

Acknowledgements

The authors are grateful to Mrs. Jacqueline Levesque and Mrs. Lise Fiset from the Cécile-Rouleau Library for their contribution to this project. They also wish to thank Mr. André Thibault for his contribution to the review of the literature in regards to the correlation between sleep curtailments and higher BMIs, as well as Mr. Jean-François Ducre-Robitaille and Mr. Bernard Caron for their constructive statistical comments. Appreciation is also expressed to Mr. John Francis Niederreiter and Mrs. Meghan Howell for the linguistic revision of this manuscript.

The authors wish to point out that the views expressed in this paper may not reflect the official policy of the Québec Ministry of Sustainable Development, Environment and Parks.

Declaration of interest

The authors report no conflicts of interest.

References

- Adams J. 2006. Socioeconomic position and sleep quantity in UK adults. *J Epidemiol Community Health* 60:267–269.
- Allen CJ, Jones NL, Killian KJ. 1984. Alveolar gas exchange during exercise: a single-breath analysis. *J Appl Physiol* 57:1704–1709.
- Andersen KL, Hart JS. 1963. Aerobic working capacity of Eskimos. *J Appl Physiol* 18:764–768.
- Andersen KL, Hermansen L. 1965. Aerobic work capacity in middle-aged Norwegian men. *J Appl Physiol* 20:432–436.
- Andersen P, Adams RP, Sjøgaard G, Thorboe A, Saltin B. 1985. Dynamic knee extension as model for study of isolated exercising muscle in humans. *J Appl Physiol* 59:1647–1653.
- Andrew GM, Guzman CA, Becklake MR. 1966. Effect of athletic training on exercise cardiac output. *J Appl Physiol* 21:603–608.
- Anton-Kuchly B, Roger P, Varenne P. 1984. Determinants of increased energy cost of submaximal exercise in obese subjects. *J Appl Physiol* 56:18–23.
- Åstrand I. 1960. Aerobic work capacity in men and women with special reference to age. *Acta Physiol Scand Suppl* 49:1–92.
- Åstrand I, Åstrand P-O, Hallbäck I, Kilbom A. 1973. Reduction in maximal oxygen uptake with age. *J Appl Physiol* 35:649–654.
- Åstrand I, Åstrand P-O, Rodahl K. 1959. Maximal heart rate during work in older men. *J Appl Physiol* 14:562–566.
- Åstrand P-O. 1952. Experimental Studies of Physical Working Capacity in Relation to Sex and Age. Kungliga Gymnastiska Central Institutet, Stockholm. Ejnar Munksengaard, Copenhagen.
- Åstrand P-O, Cuddy TE, Saltin B, Stenberg J. 1964. Cardiac output during submaximal and maximal work. *J Appl Physiol* 19:268–274.
- Åstrand P-O, Saltin B. 1961a. Oxygen uptake during the first minutes of heavy muscular exercise. *J Appl Physiol* 16:971–976.
- Åstrand P-O, Saltin B. 1961b. Maximal oxygen uptake and heart rate in various types of muscular activity. *J Appl Physiol* 16:977–981.
- Babb TG, Rodarte JR. 1993. Estimation of ventilatory capacity during submaximal exercise. *J Appl Physiol* 74:2016–2022.
- Babb TG, Viggiano R, Hurley B, Staats B, Rodarte JR. 1991. Effect of mild-to-moderate airflow limitation on exercise capacity. *J Appl Physiol* 70:223–230.
- Bachofen H, Hobi HJ, Scherrer M. 1973. Alveolar-arterial N_2 gradients at rest and during exercise in healthy men of different ages. *J Appl Physiol* 34:137–142.
- Baker SP, Hitchcock FA. 1957. Immediate effects of inhalation of 100% oxygen at one atmosphere on ventilation volume, carbon dioxide output, oxygen consumption and respiratory rate in man. *J Appl Physiol* 10:363–366.
- Bebout DE, Story D, Roca J, Hogan MC, Poole DC, Gonzalez-Camarena R, Ueno O, Haab P, Wagner PD. 1989. Effects of altitude acclimatization on pulmonary gas exchange during exercise. *J Appl Physiol* 67:2286–2295.
- Becklake MR, Frank H, Dagenais GR, Ostiguy GL, Guzman CA. 1965. Influence of age and sex on exercise cardiac output. *J Appl Physiol* 20:938–947.
- Becklake MR, Varvis CJ, Pengelly LD, Kenning S, McGregor M, Bates DV. 1962. Measurement of pulmonary blood flow during exercise using nitrous oxide. *J Appl Physiol* 17:579–586.
- Beebe DW, Lewin D, Zeller M, McCabe M, MacLeod K, Daniels SR, Amin R. 2006. Sleep in overweight adolescents: shorter sleep, poorer sleep quality, sleepiness, and sleep-disorder breathing. *J Pediatr Psychol* 104:1–11.
- Benedict FG, Carpenter TM. 1910. The Metabolism and Energy Transformations of Healthy Man during Rest. Carnegie Institute Washington Publication No. 126. The Lord Baltimore Press, Baltimore, MD.
- Bernstein MS, Costanza MC, Morabia A. 2001. Physical activity of urban adults: a general population survey in Geneva. *Soz Präventivmed* 46:49–59.
- Bessard T, Schutz Y, Jéquier E. 1983. Energy expenditure and postprandial thermogenesis in obese women before and after weight loss. *Am J Clin Nutr* 38:680–693.
- Bjorvatn B, Sagen IM, Øyane N, Waage S, Pallesen S, Ursin R. 2007. The association between sleep duration, body mass index and metabolic measures in the Hordaland Health Study. *J Sleep Res* 16:66–76.
- Blackie SP, Fairbairn MS, McElvaney NG, Wilcox PG, Morrison NJ, Pardy RL. 1991. Normal values and ranges for ventilation and breathing pattern at maximal exercise. *Chest* 100:136–142.
- Bluck LJC. 2008. Doubly labeled water for the measurements of total daily energy expenditure in man—progress and applications in the last decade. *British Nutrition Foundation. Nutr Bull* 33:80–90.
- Bollella MC, Spark A, Boccia LA, Nicklas TA, Pittman BP, Williams CL. 1999. Nutrient intake of Head Start children: home vs. school. *J Am Coll Nutr* 18:108–114.
- Bosy-Westphal A, Reinecke U, Schlörke T, Illner K, Kutzner D, Heller M, Müller MJ. 2004. Effect of organ and tissue masses on resting energy expenditure in underweight, normal weight and obese adults. *Int J Obes Relat Metab Disord* 28:72–79.
- Bransby ER, Fothergill JE. 1954. The diets of young children. *Br J Nutr* 8:195–204.

- Brault-Dubuc M, Mongeau E. 1989. Energy intake of Montreal school-age children. *J Can Diet Assoc* 50:107-112.
- Braun-Fahrlander C, Vuille JC, Sennhauser FH, Neu U, Künzle T, Grize L, Gassner M, Minder C, Schindler C, Varonier HS, Wüthrich B. 1997. Respiratory health and long-term exposure to air pollutants in Swiss schoolchildren. SCARPOL Team. Swiss Study on Childhood Allergy and Respiratory Symptoms with Respect to Air Pollution, Climate and Pollen. *Am J Respir Crit Care Med* 155:1042-1049.
- Brochu P, Ducré-Robitaille J-F, and Brodeur J. 2006a. Physiological daily inhalation rates for free-living individuals aged 1 month to 96 years, using data from doubly labeled water measurements: a proposal for air quality criteria, standard calculations and health risk assessment. *Hum Ecol Risk Assess* 12:675-701.
- Brochu P, Ducré-Robitaille J-F, Brodeur J. 2006b. Physiological daily inhalation rates for free-living pregnant and lactating adolescents and women aged 11 to 55 years, using data from doubly labeled water measurements for use in health risk assessment. *Hum Ecol Risk Assess* 12:702-735.
- Brochu P, Ducré-Robitaille J-F, Brodeur J. 2006c. Physiological daily inhalation rates for free-living individuals aged 2.6 months to 96 years based on doubly labeled water measurements: comparison with time-activity-ventilation and metabolic energy conversion estimates. *Hum Ecol Risk Assess* 12:736-761.
- Brouha L, Smith PE, De Lanne R, Maxfield ME. 1960. Physiological reactions of men and women during muscular activity and recovery in various environments. *J Appl Physiol* 16:133-140.
- Buchfuhrer MJ, Hansen JE, Robinson TE, Sue DY, Wasserman K, Whipp BJ. 1983. Optimizing the exercise protocol for cardiopulmonary assessment. *J Appl Physiol* 55:1558-1564.
- Bursztein S, Elwyn D, Askanazi J, Kinney J. 1989. Energy Metabolism, Indirect Calorimetry, and Nutrition. Williams & Wilkins, Baltimore.
- Buskirk ER, Thompson RH, Moore R, Whedon GD. 1960. Human energy expenditure studies in the national institute of arthritis and metabolic diseases metabolic chamber. *Am J Clin Nutr* 8:602-613.
- Butte NF, Calloway DH. 1981. Evaluation of lactational performance of Navajo women. *Am J Clin Nutr* 34:2210-2215.
- Butte NF, Garza C, Smith EO, Nichols BL. 1984. Human milk intake and growth in exclusively breast-fed infants. *J Pediatr* 104:187-195.
- Butte NF, Wong WW, Treuth MS, Ellis KJ, O'Brian Smith E. 2004. Energy requirements during pregnancy based on total energy expenditure and energy deposition. *Am J Clin Nutr* 79:1078-1087.
- Cade WT, Nabar SR, Keyser RE. 2004. Reproducibility of the exponential rise technique of CO₂ rebreathing for measuring P(v)CO₂ and C(v)CO₂ to non-invasively estimate cardiac output during incremental, maximal treadmill exercise. *Eur J Appl Physiol* 91:669-676.
- Caiozzo VJ, Davis JA, Berriman DJ, Vandagriff RB, Prietto CA. 1987. Effect of high-intensity exercise on the VE-VCO₂ relationship. *J Appl Physiol* 62:1460-1464.
- Cander L, Hanowell EG. 1963. Effects of fever on pulmonary diffusing capacity and pulmonary mechanics in man. *J Appl Physiol* 18:1065-1070.
- Capderou A, Douguet D, Losay J, Zelter M. 1997. Comparison of indirect calorimetry and thermodilution cardiac output measurement in children. *Am J Respir Crit Care Med* 155:1930-1934.
- Carrington MJ, Barbieri R, Colrain IM, Crowley KE, Kim Y, Trinder J. 2005. Changes in cardiovascular function during the sleep onset period in young adults. *J Appl Physiol* 98:468-476.
- Casaburi R, Whipp BJ, Wasserman K, Beaver WL, Koyal SN. 1977. Ventilatory and gas exchange dynamics in response to sinusoidal work. *J Appl Physiol* 42:300-301.
- Cizza G, Skarulis M, Mignot E. 2005. A link between short sleep and obesity: building the evidence for causation. *Sleep* 28:1217-1220.
- Cohn JE, Carroll DG, Armstrong BW, Shepard RH, Riley RL. 1954. Maximal diffusing capacity of the lung in normal male subjects of different ages. *J Appl Physiol* 6:588-597.
- Colrain IM, Trinder J, Fraser G, Wilson GV. 1987. Ventilation during sleep onset. *J Appl Physiol* 63:2067-2074.
- Cook CD, Cherry RB, O'Brien D, Karlberg P, Smith CA. 1955. Studies of respiratory physiology in the newborn infant. I. Observations on normal premature and full-term infants. *J Clin Invest* 34:975-982.
- Costill DL, Sparks K, Gregor R, Turner C. 1971. Muscle glycogen utilization during exhaustive running. *J Appl Physiol* 31:353-356.
- Craig FN. 1955. Pulmonary ventilation during exercise and inhalation of carbon dioxide. *J Appl Physiol* 7:467-471.
- Cunningham DA, Paterson DH, Koval JJ, St Croix CM. 1997. A model of oxygen transport capacity changes for independently living older men and women. *Can J Appl Physiol* 22:439-453.
- Damato AN, Galante JG, Smith WM. 1966. Hemodynamic response to treadmill exercise in normal subjects. *J Appl Physiol* 21:959-966.
- Davies CT, Godfrey S, Light M, Sargeant AJ, Zeidifard E. 1975. Cardiopulmonary responses to exercise in obese girls and young women. *J Appl Physiol* 38:373-376.
- Davis JA, Frank MH, Whipp BJ, Wasserman K. 1979. Anaerobic threshold alterations caused by endurance training in middle-aged men. *J Appl Physiol* 46:1039-1046.
- De Bock K, Richter EA, Russell AP, Eijnde BO, Derave W, Ramaekers M, Koninckx E, Léger B, Verhaeghe J, Hespel P. 2005. Exercise in the fasted state facilitates fibre type-specific intramyocellular lipid breakdown and stimulates glycogen resynthesis in humans. *J Physiol (Lond)* 564: 649-660.
- Delany JP, Schoeller DA, Hoyt RW. 1988. Use of doubly-labelled water to measure energy expenditure of special operations soldiers during a four week field training exercise FTX. *Fed Am Soc Exp Biol J* 2:5399.
- Dixon RW Jr, Faulkner JA. 1971. Cardiac outputs during maximum effort running and swimming. *J Appl Physiol* 30:653-656.
- Donevan RE, Anderson NM, Sekelj P, Papp O, McGregor M. 1962. Influence of voluntary hyperventilation on cardiac output. *J Appl Physiol* 17:487-491.
- Drinkwater BL, Horvath SM, Wells CL. 1975. Aerobic power of females, ages 10 to 68. *J Gerontol* 30:385-394.
- Durnin JV, Brockway JM, Whitcher HW. 1960. Effects of a short period of training of varying severity on some measurements of physical fitness. *J Appl Physiol* 15:161-165.
- Ehlersam RE, Heigenhauser GJ, Jones NL. 1982. Effect of respiratory acidosis on metabolism in exercise. *J Appl Physiol* 53:63-69.
- Eisenmann JC, Ekkekakis P, Holmes M. 2006. Sleep duration and overweight among Australian children and adolescents. *Acta Paediatr* 95:956-963.
- Eklom B. 1969. Effect of physical training in adolescent boys. *J Appl Physiol* 27:350-355.
- Eklom B, Astrand PO, Saltin B, Stenberg J, Wallström B. 1968. Effect of training on circulatory response to exercise. *J Appl Physiol* 24:518-528.
- Eldridge MW, Dempsey JA, Haverkamp HC, Lovering AT, Hokanson JS. 2004. Exercise-induced intrapulmonary arteriovenous shunting in healthy humans. *J Appl Physiol* 97:797-805.
- Elia M. 1997. Tissue distribution and energetics in weight loss and undernutrition. In: Physiology. Stress and Malnutrition: Functional Correlates, Nutritional Intervention, JM Kenney, HN Tucker, Eds., pp. 383-411. Lippincott-Raven Publishers, New York.
- Emirgil C, Sobol BJ, Campodonico S, Herbert WH, Mechkati R. 1967. Pulmonary circulation in the aged. *J Appl Physiol* 23:631-640.
- Eriksson BO, Grimby G, Saltin B. 1971. Cardiac output and arterial blood gases during exercise in pubertal boys. *J Appl Physiol* 31:348-352.
- Federal Register Notices. 1980. November 28. 45:79318-79379.
- Feller W. 1945. The fundamental limit theorems in probability. *Bull Amer Math Soc* 51:800-832.
- Ferrannini E. 1988. The theoretical bases of indirect calorimetry: a review. *Metab Clin Exp* 37:287-301.
- Flandrois R, Grandmontagne M, Mayet M-H, Favier R, Frutuso J. 1982. La consommation maximale d'oxygène chez le jeune français, sa variation avec l'âge, le sexe et l'entraînement. *J Physiol Paris* 78:186-194.
- Fohlin L, Freyschuss U, Bjarke B, Davies CT, Thorén C. 1978. Function and dimensions of the circulatory system in anorexia nervosa. *Acta Paediatr Scand* 67:11-16.
- Freudenheim JL, Krogh V, D'Amicis A, Scaccini C, Sette S, Ferro-Luzzi A, Trevisan M. 1993. Food sources of nutrients in the diet of elderly Italians: I. Macronutrients and lipids. *Int J Epidemiol* 22:855-868.

- Frick MH, Somer T. 1964. Base-line effects on response of stroke volume to leg exercise in the supine position. *J Appl Physiol* 19:639-643.
- Frøeb HF. 1962. Stimulation of ventilation in emphysema by passively induced body motion. *J Appl Physiol* 17:771-774.
- Frostell C, Pande JN, Hedenstierna G. 1983. Effects of high-frequency breathing on pulmonary ventilation and gas exchange. *J Appl Physiol* 55:1854-1861.
- Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB. 2005. Inadequate sleep as a risk factor for obesity: analyses of the NHANES I. *Sleep* 28:1289-1296.
- Garby L, Kurzer MS, Lammert O, Nielsen E. 1987. Energy expenditure during sleep in men and women: evaporative and sensible heat losses. *Hum Nutr: Clin Nutr* 41C:225-233.
- Gibney ER, Murgatroyd P, Wright A, Jebb S, Elia M. 2003. Measurement of total energy expenditure in grossly obese women: comparison of the bicarbonate-urea method with whole-body calorimetry and free-living doubly labelled water. *Int J Obes Relat Metab Disord* 27:641-647.
- Gibson S, Lambert J, Neate D. 2004. Association between weight status, physical activity, and consumption of biscuits, cake and confectionery among young people in Britain. *Nutr Bull* 29:301-309.
- Gisolfo J, Wilders R, Immink RV, van Lieshout JJ, Karemaker JM. 2003. Tidal volume, cardiac output and functional residual capacity determine end-tidal CO₂ transient during standing up in humans. *J Physiol (Lond)* 554:579-590.
- Godfrey S, Davies CT, Wozniak E, Barnes CA. 1971. Cardio-respiratory response to exercise in normal children. *Clin Sci* 40:419-431.
- Gore CJ, Hahn AG, Scroop GC, Watson DB, Norton KI, Wood RJ, Campbell DP, Emonson DL. 1996. Increased arterial desaturation in trained cyclists during maximal exercise at 580 m altitude. *J Appl Physiol* 80:2204-2210.
- Gross JS. 1983. Growth and biochemical response of preterm infants fed human milk or modified infant formula. *N Engl J Med* 308:237-241.
- Guyton AC. 1991. Textbook of Medical Physiology, 8th ed. W.B. Saunders Company. Harcourt Brace Jovanovich Inc., Philadelphia, PA.
- Hagberg JM, Allen WK, Seals DR, Hurley BF, Ehsani AA, Holloszy JO. 1985. A hemodynamic comparison of young and older endurance athletes during exercise. *J Appl Physiol* 58:2041-2046.
- Hagberg JM, Yerg JE 2nd, Seals DR. 1988. Pulmonary function in young and older athletes and untrained men. *J Appl Physiol* 65:101-105.
- Hanson JS. 1973. Exercise responses following production of experimental obesity. *J Appl Physiol* 35:587-591.
- Harms CA, McClaran SR, Nিকেle GA, Pegelow DE, Nelson WB, Dempsey JA. 1998. Exercise-induced arterial hypoxaemia in healthy young women. *J Physiol (Lond)* 507 (Pt 2):619-628.
- Hasler G, Buysse DJ, Klaghofer R, Gamma A, Ajdacic V, Eich D, Rössler W, Angst J. 2004. The association between short sleep duration and obesity in young adults: a 13-year prospective study. *Sleep* 27:661-666.
- Health Canada. 1996. Canadian Environmental Protection Act. Supporting Documentation: Health-Based Tolerable Daily Intakes/Concentrations and Tumourigenic Doses/Concentrations for Priority Substances. Unedited Version. Ottawa, ON, CA.
- Heath GW, Hagberg JM, Ehsani AA, Holloszy JO. 1981. A physiological comparison of young and older endurance athletes. *J Appl Physiol* 51:634-640.
- Heigenhauser GJ, Sutton JR, Jones NL. 1983. Effect of glycogen depletion on the ventilatory response to exercise. *J Appl Physiol* 54:470-474.
- Hermansen L, Andersen KL. 1965. Aerobic work capacity in young Norwegian men and women. *J Appl Physiol* 20:425-431.
- Hermansen L, Ekblom B, Saltin B. 1970. Cardiac output during submaximal and maximal treadmill and bicycle exercise. *J Appl Physiol* 29:82-86.
- Hermansen L, Saltin B. 1969. Oxygen uptake during maximal treadmill and bicycle exercise. *J Appl Physiol* 26:31-37.
- Hitchcock NE, Owles EN, Gracey M, Gilmour A. 1984. Nutrition of healthy children in the second and third years of life. *J Food Nutr* 41:13-16.
- Hoffmans MD, Obermann-de Boer GL, Florack EI, van Kampen-Donker M, Kromhout D. 1986. Energy, nutrient and food intake during infancy and early childhood. The Leiden Preschool Children Study. *Hum Nutr Appl Nutr* 40:421-430.
- Holmér I. 1972. Oxygen uptake during swimming in man. *J Appl Physiol* 33:502-509.
- Hudgel DW, Devadatta P, Hamilton H. 1993. Pattern of breathing and upper airway mechanics during wakefulness and sleep in healthy elderly humans. *J Appl Physiol* 74:2198-2204.
- Hunter GR, Weinsier RL, Darnell BE, Zuckerman PA, Goran MI. 2000. Racial differences in energy expenditure and aerobic fitness in premenopausal women. *Am J Clin Nutr* 71:500-506.
- Iglowstein I, Jenni OG, Molinari L, Largo RH. 2003. Sleep duration from infancy to adolescence: reference values and generational trends. *Pediatrics* 111:302-307.
- Inbar O, Bar-Or O, Dotan R, Gutin B. 1981. Conditioning versus exercise in heat as methods for acclimatizing 8- to 10-yr-old boys to dry heat. *J Appl Physiol* 50:406-411.
- Institute of Medicine (IOM). 2002. Appendix I: doubly labeled water data used to predict energy expenditure. In: Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients), pp. 1104-1202. Food and Nutrition Board. National Academies Press, Washington, DC. Available at: <http://books.nap.edu/books/0309085373/html/index.html>
- International Dietary Energy Consultancy Group (IDECG). 1990. The Doubly-Labelled Water Method for Measuring Energy Expenditure: A Consensus Report by the IDECG working group. Technical recommendation for use in humans. NAHRES-4, IAEA. Prentice AM, Ed. Vienna, Austria. Available at: <http://www.unu.edu/unupress/food2/UID05E/UID05E00.HTM>
- Jenner DA, English DR, Vandongen R, Beilin LJ, Armstrong BK, Miller MR, Dunbar D. 1988. Diet and blood pressure in 9-year-old Australian children. *Am J Clin Nutr* 47:1052-1059.
- Johnson RL Jr, Spicer WS, Bishop JM, Forster RE. 1960. Pulmonary capillary blood volume, flow and diffusing capacity during exercise. *J Appl Physiol* 15:893-902.
- Jones NL, Robertson DG, Kane JW. 1979. Difference between end-tidal and arterial PCO₂ in exercise. *J Appl Physiol* 47:954-960.
- Jones PJ, Winthrop AL, Schoeller DA, Filler RM, Swyer PR, Smith J, Heim T. 1988. Evaluation of doubly labeled water for measuring energy expenditure during changing nutrition. *Am J Clin Nutr* 47:799-804.
- Jones WB, Finchum RN, Russell RO Jr, Reeves TJ. 1970. Transient cardiac output response to multiple levels of supine exercise. *J Appl Physiol* 28:183-189.
- Joyner MJ, Freund BJ, Jilka SM, Hetrick GA, Martinez E, Ewy GA, Wilmore JH. 1986. Effects of β -blockade on exercise capacity of trained and untrained men: a hemodynamic comparison. *J Appl Physiol* 60:1429-1434.
- Kanstrup IL, Ekblom B. 1978. Influence of age and physical activity on central hemodynamics and lung function in active adults. *J Appl Physiol* 45:709-717.
- Karlsson J, Astrand PO, Ekblom B. 1967. Training of the oxygen transport system in man. *J Appl Physiol* 22:1061-1065.
- Kasch FW, Boyer JL, Van Camp S, Nettl F, Verity LS, Wallace JP. 1995. Cardiovascular changes with age and exercise. A 28-year longitudinal study. *Scand J Med Sci Sports* 5:147-151.
- Kastello GM, Sothmann MS, Murthy VS. 1993. Young and old subjects matched for aerobic capacity have similar noradrenergic responses to exercise. *J Appl Physiol* 74:49-54.
- Katayama K, Sato Y, Morotome Y, Shima N, Ishida K, Mori S, Miyamura M. 1999. Ventilatory chemosensitive adaptations to intermittent hypoxic exposure with endurance training and detraining. *J Appl Physiol* 86:1805-1811.
- Kearney JT, Stull GA, Ewing JL Jr, Strein JW. 1976. Cardiorespiratory responses of sedentary college women as a function of training intensity. *J Appl Physiol* 41:822-825.

- Knuiman JT, Westenbrink S, van der Heyden L, West CE, Burema J, de Boer J, Hautvast JG, Räsänen L, Virkkunen L, Viikari J. 1983. Determinants of total and high density lipoprotein cholesterol in boys from Finland, The Netherlands, Italy, the Philippines and Ghana with special reference to diet. *Hum Nutr Clin Nutr* 37:237-254.
- Knutson KL, Lauderdale DS. 2007. Sleep duration and overweight in adolescents: self-reported sleep hours versus time diaries. *Pediatrics* 119:e1056-e1062.
- Knuttgen HG. 1967. Aerobic capacity of adolescents. *J Appl Physiol* 22:655-658.
- Kobayashi K, Kitamura K, Miura M, Sodeyama H, Murase Y, Miyashita M, Matsui H. 1978. Aerobic power as related to body growth and training in Japanese boys: a longitudinal study. *J Appl Physiol* 44:666-672.
- Koch G, Eriksson BO. 1973. Effect of physical training on anatomical R-L shunt at rest and pulmonary diffusing capacity during near-maximal exercise in boys 11-13 years old. *Scand J Clin Lab Invest* 31:95-103.
- Koçoglu G, Ozdemir L, Sümer H, Demir DA, Cetinkaya S, Polat HH. 2003. Prevalence of obesity among 11-14 years old students in Sivas-Turkey. *Pakistan J Nutr* 2:292-295.
- Kohatsu ND, Tsai R, Young T, Vangilder R, Burmeister LE, Stromquist AM, Merchant JA. 2006. Sleep duration and body mass index in a rural population. *Arch Intern Med* 166:1701-1705.
- Kohrt WM, Malley MT, Coggan AR, Spina RJ, Ogawa T, Ehsani AA, Bourey RE, Martin WH 3rd, Holloszy JO. 1991. Effects of gender, age, and fitness level on response of $\dot{V}O_{2max}$ to training in 60-71 yr olds. *J Appl Physiol* 71:2004-2011.
- Kripke DE, Garfinkel L, Wingard DL, Klauber MR, Marler MR. 2002. Mortality associated with sleep duration and insomnia. *Arch Gen Psychiatry* 59:131-136.
- Krone RJ, Goldbarg AN, Balkoura M, Schuessler R, Resnekov L. 1972. Effects of cigarette smoking at rest and during exercise. II. Role of venous return. *J Appl Physiol* 32:745-748.
- Layton DW. 1993. Metabolically consistent breathing rates for use in dose assessments. *Health Phys* 64:23-36.
- Lees MH, Way RC, Ross BB. 1967. Ventilation and respiratory gas transfer of infants with increased pulmonary blood flow. *Pediatrics* 40:259-271.
- Leung M, Yeung DL, Pennell MD, Hall J. 1984. Dietary intakes of preschoolers. *J Am Diet Assoc* 84:551-554.
- Lewis SF, Taylor WF, Graham RM, Pettinger WA, Schutte JE, Blomqvist CG. 1983. Cardiovascular responses to exercise as functions of absolute and relative work load. *J Appl Physiol* 54:1314-1323.
- Liu S, Krewski D, Shi Y, Chen Y, Burnett RT. 2003. Association between gaseous ambient air pollutants and adverse pregnancy outcomes in Vancouver, Canada. *Environ Health Perspect* 111:1773-1778.
- Loftin M, Sothorn M, Trosclair L, O'Hanlon A, Miller J, Udall J. 2001. Scaling $\dot{V}O_2$ peak in obese and non-obese girls. *Obes Res* 9:290-296.
- Makrides L, Heigenhauser GJ, Jones NL. 1990. High-intensity endurance training in 20- to 30- and 60- to 70-yr-old healthy men. *J Appl Physiol* 69:1792-1798.
- Malcom A, Holliday MD. 1971. Metabolic rate and organ size during growth from infancy to maturity and during late gestation and early infancy. *Pediatrics* 47:169-179.
- Malmberg R. 1966. Pulmonary gas exchange at exercise and different body postures in man. *Scand J Respir Dis* 47:92-102.
- Margarey A, Boulton JC. 1984. Nutritional studies during childhood: IV Energy and nutrient intake at age 4. *Aust Paediatr J* 20:187-194.
- Marti B, Howald H. 1990. Long-term effects of physical training on aerobic capacity: controlled study of former elite athletes. *J Appl Physiol* 69:1451-1459.
- Martin B, Heintzelman M, Chen HI. 1982. Exercise performance after ventilatory work. *J Appl Physiol* 52:1581-1585.
- Marven SS, Smith CM, Claxton D, Chapman J, Davies HA, Primhak RA, Powell CV. 1998. Pulmonary function, exercise performance, and growth in survivors of congenital diaphragmatic hernia. *Arch Dis Child* 78:137-142.
- Mason RJ, Broaddus VC, Murray JE, Nadel JA. 2005. *Murray and Nadel's Textbook of Respiratory Medicine*, 4th ed. Saunders, Philadelphia, PA.
- McClaran SR, Babcock MA, Pegelow DE, Reddan WG, Dempsey JA. 1995. Longitudinal effects of aging on lung function at rest and exercise in healthy active fit elderly adults. *J Appl Physiol* 78:1957-1968.
- McLean JA, Tobin G. 1987. *Animal and Human Calorimetry*. Cambridge University Press, Cambridge, MA.
- Michael ED Jr, Horvath SM. 1965. Physical work capacity of college women. *J Appl Physiol* 20:263-266.
- Miyamoto Y, Hiura T, Tamura T, Nakamura T, Higuchi J, Mikami T. 1982. Dynamics of cardiac, respiratory, and metabolic function in men in response to step workload. *J Appl Physiol* 52:1198-1208.
- Miyamoto Y, Niizeki K, Kawahara K, Doi K. 1989. Cardiodynamic factors affecting hyperpnea during steady-state exercise in man. *Jpn J Physiol* 39:411-420.
- Miyamura M, Honda Y. 1972. Oxygen intake and cardiac output during maximal treadmill and bicycle exercise. *J Appl Physiol* 32:185-188.
- Morisson JA, Larsen R, Glatfelter L, Boggs D, Burton K, Smith C, Kelly K, Mellies MJ, Khoury P, Glueck CJ. 1980. Nutrient intake: relationships with lipids and lipoproteins in 6-19-year-old children—the Princeton School District study. *Metab Clin Exp* 29:133-140.
- Morrell MJ, Harty HR, Adams L, Guz A. 1995. Changes in total pulmonary resistance and PCO_2 between wakefulness and sleep in normal human subjects. *J Appl Physiol* 78:1339-1349.
- Mosteller RD. 1987. Simplified calculation of body-surface area. *N Engl J Med* 317:1098.
- Mostyn EM, Helle S, Gee JB, Bentivoglio LG, Bates DV. 1963. Pulmonary diffusing capacity of athletes. *J Appl Physiol* 18:687-695.
- Müller MJ, von zur Mühlen A, Lautz HU, Schmidt FW, Daiber M, Hürter P. 1989. Energy expenditure in children with type I diabetes: evidence for increased thermogenesis. *BMJ* 299:487-491.
- Murphy TM, Clark WH, Buckingham IP, Young WA. 1969. Respiratory gas exchange in exercise during helium-oxygen breathing. *J Appl Physiol* 26:303-307.
- Naimark A, Wasserman K, McIlroy MB. 1964. Continuous measurement of ventilatory exchange ratio during exercise. *J Appl Physiol* 19:644-652.
- Narasinga RBS, Susheela TP, Nadamuni Naidu A, Menon K. 1982. Energy intake of well-to-do preschool children in India. *Indian J Med Res* 77:62-72.
- Neder JA, Dal Corso S, Malaguti C, Reis S, De Fuccio MB, Schmidt H, Fuld JP, Nery LE. 2003. The pattern and timing of breathing during incremental exercise: a normative study. *Eur Respir J* 21:530-538.
- Neiderud J, Philip I, Sjölin S. 1992. Greek immigrant children in southern Sweden in comparison with Greek and Swedish children. III. Energy and nutrient intake. *Acta Paediatr* 81:430-435.
- Nelson M, Naismith DJ, Burley V, Gatenby S, Geddes N. 1990. Nutrient intakes, vitamin-mineral supplementation, and intelligence in British schoolchildren. *Br J Nutr* 64:13-22.
- Nelson NM, Prod'Hom LS, Cherry RB, Lipsitz PJ, Smith CA. 1962. Pulmonary function in the newborn infant. I. Methods: ventilation and gaseous metabolism. *Pediatrics* 30:963-974.
- Nery LE, Wasserman K, Andrews JD, Huntsman DJ, Hansen JE, Whipp BJ. 1982. Ventilatory and gas exchange kinetics during exercise in chronic airways obstruction. *J Appl Physiol* 53:1594-1602.
- Newman F, Smalley BF, Thomson ML. 1962. Effect of exercise, body and lung size on CO diffusion in athletes and nonathletes. *J Appl Physiol* 17:649-655.
- Nottin S, Vinet A, Lecoq AM, Guenon P, Obert P. 2000. [Study of the reproducibility of cardiac output measurement during exercise in pre-pubertal children by Doppler echocardiography and CO_2 inhalation]. *Arch Mal Coeur Vaiss* 93:1297-1303.

- Nourry C, Deruelle F, Fabre C, Baquet G, Bart F, Grosbois JM, Berthoin S, Mucci P. 2005. Exercise flow-volume loops in prepubescent aerobically trained children. *J Appl Physiol* 99:1912-1921.
- Nutrition Canada (NC). 1977. Food Consumption Patterns Report (CA1 HW 12077F51). Bureau of Nutritional Sciences, Health Protection Branch, Department of National Health and Welfare, CA.
- Ocel JV, Miller LE, Pierson LM, Wooten DF, Hawkins BJ, Myers J, Herbert WG. 2003. Adaptation of pulmonary oxygen consumption slow component following 6 weeks of exercise training above and below the lactate threshold in untrained men. *Chest* 124:2377-2383.
- Olfert IM, Balouch J, Kleinsasser A, Knapp A, Wagner H, Wagner PD, Hopkins SR. 2004. Does gender affect human pulmonary gas exchange during exercise? *J Physiol (Lond)* 557:529-541.
- Oliveria SA, Ellison RC, Moore LL, Gillman MW, Garrahe EJ, Singer MR. 1992. Parent-child relationships in nutrient intake: the Framingham Children's Study. *Am J Clin Nutr* 56:593-598.
- Ong HY, O'Dochartaigh CS, Lovell S, Patterson VH, Wasserman K, Nicholls DP, Riley MS. 2004. Gas exchange responses to constant work-rate exercise in patients with glycogenosis type V and VII. *Am J Respir Crit Care Med* 169:1238-1244.
- Oren A, Wasserman K, Davis JA, Whipp BJ. 1981. Effect of CO₂ set point on ventilatory response to exercise. *J Appl Physiol* 51:185-189.
- Osborne MA, Schneider DA. 2005. Muscle glycogen reduction in man: relationship between surface EMG activity and oxygen uptake kinetics during heavy exercise. *Exp Physiol* 91:179-189.
- Ouellet Y, Poh SC, Becklake MR. 1969. Circulatory factors limiting maximal aerobic exercise capacity. *J Appl Physiol* 27:874-880.
- Pao EM, Mickle SJ, Burk MC. 1985. One-day and 3-day nutrient intakes by individuals—Nationwide Food Consumption Survey findings, Spring 1977. *J Am Diet Assoc* 85:313-324.
- Patel SR, Ayas NT, Malhotra MR, White DP, Schernhammer ES, Speizer FE, Stampfer MJ, Hu FB. 2004. A prospective study of sleep duration and mortality risk in women. *Sleep* 27:440-444.
- Patel SR, Malhotra A, White DP, Gottlieb DJ, Hu FB. 2006. Association between reduced sleep and weight gain in women. *Am J Epidemiol* 164:947-954.
- Payne JA, Belton NR. 1992. Nutrient intake and growth in preschool children. I. Comparison of energy intake and source of energy with growth. *J Hum Nutr Diet* 5:287-298.
- Pellegrino R, Villosio C, Milanese U, Garelli G, Rodarte JR, Brusasco V. 1999. Breathing during exercise in subjects with mild-to-moderate airflow obstruction: effects of physical training. *J Appl Physiol* 87:1697-1704.
- Pernow B, Saltin B. 1971. Availability of substrates and capacity for prolonged heavy exercise in man. *J Appl Physiol* 31:416-422.
- Petersson J, Sánchez-Crespo A, Rohdin M, Montmerle S, Nyén S, Jacobsson H, Larsson SA, Lindahl SG, Linnarsson D, Glenney RW, Mure M. 2004. Physiological evaluation of a new quantitative SPECT method measuring regional ventilation and perfusion. *J Appl Physiol* 96:1127-1136.
- Pierce AK, Luterma D, Loudermilk J, Blomqvist G, Johnson RL Jr. 1968. Exercise ventilatory patterns in normal subjects and patients with airway obstruction. *J Appl Physiol* 25:249-254.
- Pimentel AE, Gentile CL, Tanaka H, Seals DR, Gates PE. 2003. Greater rate of decline in maximal aerobic capacity with age in endurance-trained than in sedentary men. *J Appl Physiol* 94:2406-2413.
- Plowman ML, Drinkwater BK, Horvath SM. 1979. Age and aerobic power in women: a longitudinal study. *J Gerontol* 34:512-520.
- Podolsky A, Eldridge MW, Richardson RS, Knight DR, Johnson EC, Hopkins SR, Johnson DH, Michimata H, Grassi B, Feiner J, Kurdak SS, Bickler PE, Severinghaus JW, Wagner PD. 1996. Exercise-induced VA/Q inequality in subjects with prior high-altitude pulmonary edema. *J Appl Physiol* 81:922-932.
- Polgar G, Weng TR. 1979. The functional development of the respiratory system from the period of gestation to adulthood. *Am Rev Respir Dis* 120:625-695.
- Pollock ML, Miller HS Jr, Janeway R, Linnerud AC, Robertson B, Valentino R. 1971. Effects of walking on body composition and cardiovascular function of middle-aged man. *J Appl Physiol* 30:126-130.
- Poole JG, Lawrenson L, Kim J, Brown C, Richardson RS. 2002. Vascular and metabolic response to cycle exercise in sedentary humans: effect of age. *Am J Physiol Heart Circ Physiol* 284:H1251-H1259.
- Proctor DN, Beck KC. 1996. Delay time adjustments to minimize errors in breath-by-breath measurement of VO₂ during exercise. *J Appl Physiol* 81:2495-2499.
- Pugh LGCE. 1964. Cardiac output in muscular exercise at 5,800 m (19,000 ft). *J Appl Physiol* 19:441-447.
- Putman CT, Jones NL, Hultman E, Hollidge-Horvat MG, Bonen A, McConachie DR, Heigenhauser GJ. 1998. Effects of short-term submaximal training in humans on muscle metabolism in exercise. *Am J Physiol* 275:E132-E139.
- Raine JM, Bishop JM. 1963. A difference in O₂ tension and physiological dead space in normal man. *J Appl Physiol* 18:284-288.
- Räsänen L, Ahola M, Kara R, Uhari M. 1985. Atherosclerosis precursors in Finnish children and adolescents. VIII. Food consumption and nutrient intakes. *Acta Paediatr Scand Suppl* 318:135-153.
- Räsänen L, Laitinen S, Stirkkinen R, Kimppa S, Viikari J, Uhari M, Pesonen E, Salo M, Akerblom HK. 1991. Composition of the diet of young Finns in 1986. *Ann Med* 23:73-80.
- Räsänen L, Ylönen K. 1992. Food consumption and nutrient intake of one- to two-year-old Finnish children. *Acta Paediatr* 81:7-11.
- Ravussin E, Burnand B, Schutz Y, Jéquier E. 1985. Energy expenditure before and during energy restriction in obese patients. *Am J Clin Nutr* 41:753-759.
- Razanamahefa L, Lafay L, Oseredczuk M, Thiébaud A, Laloux L, Gerber M, Astorg P, Berta J-L. 2005. Consommation lipidique de la population française et qualité des données de composition des principaux groupes d'aliments vecteurs. *Bull Cancer* 92:647-657.
- Reeves JT, Grover RF, Blount SG Jr, Filley GF. 1961. Cardiac output response to standing and treadmill walking. *J Appl Physiol* 16:283-288.
- Reichman B, Chessex P, Putet G, Verellen G, Smith JM, Heim T, Swyer PR. 1981. Diet, fat accretion, and growth in premature infants. *N Engl J Med* 305:1495-1500.
- Renwick AG. 2000. The use of safety or uncertainty factors in the setting of acute reference doses. *Food Addit Contam* 17:627-635.
- Riboli E, Elmstahl S, Saracci R, Gullberg B, Lindgärde F. 1997. The Malmö Food Study: validity of two dietary assessment methods for measuring nutrient intake. *Int J Epidemiol* 26 (Suppl 1):S161-S173.
- Rice JA. 1995. Mathematical Statistics and Data Analysis, 2nd ed. Duxbury Press, Belmont, CA.
- Riddell MC, Bar-Or O, Wilk B, Parolin ML, Heigenhauser GJ. 2001. Substrate utilization during exercise with glucose and glucose plus fructose ingestion in boys ages 10-14 yr. *J Appl Physiol* 90:903-911.
- Robinson S. 1938. Experimental studies of physical fitness in relation to age. *Arbeits Physiol* 10:251-323.
- Roca J, Agustí AG, Alonso A, Poole DC, Viegas C, Barbera JA, Rodríguez-Roisin R, Ferrer A, Wagner PD. 1992. Effects of training on muscle O₂ transport at VO_{2max}. *J Appl Physiol* 73:1067-1076.
- Rogers MA, Hagberg JM, Martin WH 3rd, Ehsani AA, Holloszy JO. 1990. Decline in VO_{2max} with aging in master athletes and sedentary men. *J Appl Physiol* 68:2195-2199.
- Rowland TW, Boyajian A. 1995. Aerobic response to endurance exercise training in children. *Pediatrics* 96:654-658.
- Rowland TW, Cunningham LN. 1997. Development of ventilatory responses to exercise in normal white children. A longitudinal study. *Chest* 111:327-332.
- Ruxton CHS, Kirk TR, Belton NR. 1996. The contribution of specific dietary patterns to energy and nutrient intakes in 7-8-year-old Scottish schoolchildren. II Weekday lunches. *J Hum Nutr Diet* 9:15-22.
- Saltzman HA, Salzano JV. 1971. Effects of carbohydrate metabolism upon respiratory gas exchange in normal men. *J Appl Physiol* 30:228-231.
- Sargeant AJ, Rouleau MY, Sutton JR, Jones NL. 1981. Ventilation in exercise studied with circulatory occlusion. *J Appl Physiol* 50:718-723.
- Schiller BC, Casas YG, Desouza CA, Seals DR. 2001. Maximal aerobic capacity across age in healthy Hispanic and Caucasian women. *J Appl Physiol* 91:1048-1054.

- Schutz Y, Bessard T, Jéquier E. 1984. Diet-induced thermogenesis measured over a whole day in obese and nonobese women. *Am J Clin Nutr* 40:542-552.
- Segal SS, Brooks GA. 1979. Effects of glycogen depletion and work load on postexercise O_2 consumption and blood lactate. *J Appl Physiol* 47:514-521.
- Seicean A, Redline S, Seicean S, Kirchner HL, Gao Y, Sekine M, Zhu X, Storfer-Isser A. 2007. Association between short sleeping hours and overweight in adolescents: results from a US Suburban High School survey. *Sleep* 30:285-293.
- Shapiro CM, Goll CC, Cohen GR, Oswald I. 1984. Heat production during sleep. *J Appl Physiol* 56:671-677.
- Sharma JD, Saxena RK, Rastogi SK. 1977. Cardiac output of Indian men by a non-invasive method the Indirect Fick Principle. *Indian J Physiol Pharmacol* 21:347-352.
- Shepherd AP, Terpolilli BM, Steinke JM. 2007. A hand-held device to measure oxygen uptake: performance characteristics, patient selection and the propagation of its measurement error into fick cardiac output determinations. *J Invasive Cardiol* 19:113-122.
- Shiou-Liang W, Clyde W, Kostas T, Leslie B. 2005. Ingestion of high-glycemic index meal increases muscle glycogen storage at rest but augments its utilization during subsequent exercise. *J Appl Physiol* 99:707-714.
- Sidney KH, Shephard RJ. 1977. Maximum and submaximum exercise tests in men and women in the seventh, eighth, and ninth decades of life. *J Appl Physiol* 43:280-287.
- Simons-Morton DG, Hunsberger SA, Van Horn L, Barton BA, Robson AM, McMahon RP, Muhonen LE, Kwiterovich PO, Lasser NL, Kimm SY, Greenlick MR. 1997. Nutrient intake and blood pressure in the Dietary Intervention Study in Children. *Hypertension* 29:930-936.
- Sinning WE, Adrian MJ. 1968. Cardiorespiratory changes in college women due to a season of competitive basketball. *J Appl Physiol* 25:720-724.
- Skoog DA, Holler FJ, Crouch SR. 2006. Principles of Instrumental Analysis, 6th ed., pp. 1056. Brooks/Cole Publishing Company, Pacific Grove, CA.
- Spurr GB, Hutt BK, Horvath SM. 1957. Shivering, oxygen consumption and body temperatures in acute exposure of men to two different cold environments. *J Appl Physiol* 11:58-64.
- Stahlman MT, Meece NJ. 1957. Pulmonary ventilation and diffusion in the human newborn infant. *J Clin Invest* 36:1081-1091.
- Stamford BA. 1975. Maximal oxygen uptake during treadmill walking and running at various speeds. *J Appl Physiol* 39:386-389.
- Stevenson ET, Davy KP, Seals DR. 1994. Maximal aerobic capacity and total blood volume in highly trained middle-aged and older female endurance athletes. *J Appl Physiol* 77:1691-1696.
- Stickland MK, Welsh RC, Petersen SR, Tyberg JV, Anderson WD, Jones RL, Taylor DA, Bouffard M, Haykowsky MJ. 2006. Does fitness level modulate the cardiovascular hemodynamic response to exercise? *J Appl Physiol* 100:1895-1901.
- Sue DY, Wasserman K, Moricca RB, Casaburi R. 1988. Metabolic acidosis during exercise in patients with chronic obstructive pulmonary disease. Use of the V-slope method for anaerobic threshold determination. *Chest* 94:931-938.
- Tabachnik E, Muller NL, Bryan AC, Levison H. 1981. Changes in ventilation and chest wall mechanics during sleep in normal adolescents. *J Appl Physiol* 51:557-564.
- Tabakin BS, Hanson JS, Merriam TW Jr, Caldwell EJ. 1964. Hemodynamic response of normal men to graded treadmill exercise. *J Appl Physiol* 19:457-464.
- Taheri S. 2006. The link between short sleep duration and obesity: we should recommend more sleep to prevent obesity. *Arch Dis Child* 91:881-884.
- Taheri S, Lin L, Austin D, Young T, Mignot E. 2004. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* 1:e62.
- Tanaka H, Desouza CA, Jones PP, Stevenson ET, Davy KP, Seals DR. 1997. Greater rate of decline in maximal aerobic capacity with age in physically active vs. sedentary healthy women. *J Appl Physiol* 83:1947-1953.
- Tenney SM, Miller RM. 1956. Dead space ventilation in old age. *J Appl Physiol* 9:321-327.
- Thang NM, Popkin BM. 2004. Patterns of food consumption in Vietnam: effects on socioeconomic groups during an era of economic growth. *Eur J Clin Nutr* 58:145-153.
- Tokudome Y, Imaeda N, Ikeda M, Kitagawa I, Fujiwara N, Tokudome S. 1999. Foods contributing to absolute intake and variance in intake of fat, fatty acids and cholesterol in middle-aged Japanese. *J Epidemiol* 9:78-90.
- Tolbert PE, Mulholland JA, MacIntosh DL, Xu F, Daniels D, Devine OJ, Carlin BP, Klein M, Dorley J, Butler AJ, Nordenberg DE, Frumkin H, Ryan PB, White MC. 2000. Air quality and pediatric emergency room visits for asthma in Atlanta, Georgia, USA. *Am J Epidemiol* 151:798-810.
- Toner MM, Sawka MN, Levine L, Pandolf KB. 1983. Cardiorespiratory responses to exercise distributed between the upper and lower body. *J Appl Physiol* 54:1403-1407.
- Torre-Bueno JR, Wagner PD, Saltzman HA, Gale GE, Moon RE. 1985. Diffusion limitation in normal humans during exercise at sea level and simulated altitude. *J Appl Physiol* 58:989-995.
- Torun B, Davies PS, Livingstone MB, Paolisso M, Sackett R, Spurr GB. 1996. Energy requirements and dietary energy recommendations for children and adolescents 1 to 18 years old. *Eur J Clin Nutr* 50 (Suppl 1):S37-S80; discussion S80.
- Truth MS, Adolph AL, Butte NF. 1998. Energy expenditure in children predicted from heart rate and activity calibrated against respiration calorimetry. *Am J Physiol* 275:E12-E18.
- Trotter HF. 1959. An elementary proof of the central limit theorem. *Arch Math* 10:226-234.
- Turel DJ, Alexander JK. 1964. Experimental evaluation of Weir's formula for estimating metabolic rate in man. *J Appl Physiol* 19: 946-948.
- U.S. Department of Agriculture (USDA). 1984. Nutrient Intakes: Individuals in the United States, years 1977-1978, NFCS 1977-1978. Report No. I-2. U.S. Dept. of Agriculture, Human Nutrition Information Service, Washington, DC.
- U.S. Department of Health and Human Services (DHHS). 1983. Dietary Intakes Source Data: United States, 1976-1980. Publication No. (PHS) 83-1681. National Center for Health Statistics, Hyattsville, MD.
- van Engelen JGM, Prud'homme de Lodder LCH. 2007. Non-Food Products: How to Assess Children's Exposure? RIVM report 320005005/2007. Centre for Substances and Integrated Risk Assessment, National Institute of Public Health and Environmental Protection, Bilthoven. Available at: <http://rivm.openrepository.com/rivm/bitstream/10029/16474/1/320005005.pdf>
- Veicsteinas A, Samaja M, Gussoni M, Cerretelli P. 1984. Blood O_2 affinity and maximal O_2 consumption in elite bicycle racers. *J Appl Physiol* 57:52-58.
- Vogel JA, Patton JF, Mello RP, Daniels WL. 1986. An analysis of aerobic capacity in a large United States population. *J Appl Physiol* 60:494-500.
- Volianitis S, Yoshiga CC, Nissen P, Secher NH. 2003. Effect of fitness on arm vascular and metabolic responses to upper body exercise. *Am J Physiol Heart Circ Physiol* 286:H1736-H1741.
- Vorona RD, Winn MP, Babineau TW, Eng BP, Feldman HR, Ware JC. 2005. Overweight and obese patients in a primary care population report less sleep than patients with a normal body mass index. *Arch Intern Med* 165:25-30.
- Wagner PD, Gale GE, Moon RE, Torre-Bueno JR, Stolp BW, Saltzman HA. 1986. Pulmonary gas exchange in humans exercising at sea level and simulated altitude. *J Appl Physiol* 61:260-270.
- Weber G, Kartodihardjo W, Klissouras V. 1976. Growth and physical training with reference to heredity. *J Appl Physiol* 40:211-215.
- Weir JB. 1949. New methods for calculating metabolic rate with special reference to protein metabolism. *J Physiol (Lond)* 109:1-9.
- Whipp BJ, Wasserman K. 1969. Alveolar-arterial gas tension differences during graded exercise. *J Appl Physiol* 27:361-365.
- Woo J, Leung SS, Ho SC, Lam TH, Janus ED. 1998. Dietary intake and practices in the Hong Kong Chinese population. *J Epidemiol Community Health* 52:631-637.

- Woo JS, Derleth C, Stratton JR, Levy WC. 2006. The influence of age, gender, and training on exercise efficiency. *J Am Coll Cardiol* 47:1049-1057.
- World Health Organization (WHO). 2005. Chemical-Specific and Adjustment Factors for Interspecies Differences and Human Variability: Guidance Document for Use of Data in Dose/Concentration-Response Assessment. Harmonization Project Document No. 2. WHO Library Cataloguing-in-Publication Data. ISBN 92 4 154678 6. Geneva.
- Yamaji K, Miyashita M. 1977. Oxygen transport system during exhaustive exercise in Japanese boys. *Eur J Appl Physiol Occup Physiol* 36:93-99.
- Yang Q, Chen Y, Shi Y, Burnett RT, McGrail KM, Krewski D. 2003. Association between ozone and respiratory admissions among children and the elderly in Vancouver, Canada. *Inhal Toxicol* 15:1297-1308.
- Yerg JE 2nd, Seals DR, Hagberg JM, Holloszy JO. 1985. Effect of endurance exercise training on ventilatory function in older individuals. *J Appl Physiol* 58:791-794.
- Zaregarizi M, Edwards B, George K, Harrison Y, Jones H, Atkinson G. 2007. Acute changes in cardiovascular function during the onset period of daytime sleep: comparison to lying awake and standing. *J Appl Physiol* 103:1332-1338.
- Zimmerman P, Heigenhauser GJ, McCartney N, Sutton JR, Jones NL. 1982. Impaired cardiac "acceleration" at the onset of exercise in patients with coronary disease. *J Appl Physiol* 52:71-78.

Appendix

VO₂ and VE measurements reported in the following studies were used to calculate the VQ values of Table 8. Values for VO₂ and VCO₂ taken from the underlined references were also used to determine the *H* values.

Individual data from Robinson (1938), Åstrand (1952), Cohn et al. (1954), Cook et al. (1955), Craig (1955), Stahlman and Meece (1957), Åstrand et al. (1959), Åstrand and Saltin (1961a, b), Becklake et al. (1962), Donevan et al. (1962), Nelson et al. (1962), Newman et al. (1962), Andersen and Hart (1963), Cander and Hanowell (1963), Mostyn et al. (1963), Pugh (1964), Tabakin et al. (1964), Michael and Horvath (1965), Damato et al. (1966), Karlsson et al. (1967), Eklblom et al. (1968), Pierce et al. (1968), Murphy et al. (1969), Ouellet et al. (1969), Costill et al. (1971), Holmér (1972), Bachofen et al. (1973), Casaburi et al. (1977), Kobayashi et al. (1978), Jones et al. (1979), Frostell et al. (1983), Martin et al. (1982), Torre-Bueno et al. (1985), Babb and Rodarte (1993), Eldridge et al. (2004), and Ong et al. (2004).

Mean values and standard deviations were from Åstrand (1960), Brouha et al. (1960), Durnin et al. (1960), Froeb (1962), Raine and Bishop (1963), Naimark et al. (1964), Andersen and Hermansen (1965), Becklake et al. (1965), Hermansen and Andersen (1965), Andrew et al. (1966), Malmberg (1966), Knuttgen (1967), Sinning and Adrian (1968), Eklblom (1969), Hermansen and Saltin (1969), Whipp and Wasserman (1969), Dixon and Faulkner (1971), Eriksson et al. (1971), Godfrey et al. (1971), Pollock et al. (1971), Krone et al. (1972), Miyamura and Honda (1972), Åstrand et al. (1973), Hanson (1973), Koch and Eriksson (1973), Davies et al. (1975), Drinkwater et al. (1975), Stamford (1975), Kearney et al. (1976), Weber et al. (1976), Sharma et al. (1977), Sidney and Shephard (1977), Yamaji and Miyashita (1977), Fohlin et al. (1978), Kanstrup and Eklblom (1978), Davis et al. (1979), Plowman et al. (1979), Segal and Brooks (1979), Heath et al. (1981), Inbar et al. (1981), Sargeant et al. (1981), Ehram et al. (1982), Flandrois et al. (1982), Miyamoto et al. (1982), Nery et al. (1982), Zimmerman et al. (1982), Buchfuhrer et al. (1983), Heigenhauser et al. (1983), Lewis et al. (1983), Toner et al. (1983), Anton-Kuchly et al. (1984), Veicsteinas et al. (1984), Andersen et al. (1985), Hagberg et al. (1985), Yerg et al. (1985), Joyner et al. (1986), Vogel et al. (1986), Wagner et al. (1986), Caiozzo et al. (1987), Hagberg et al. (1988), Sue et al. (1988), Bebout et al. (1989), Miyamoto et al. (1989), Makrides et al. (1990), Marti and Howald (1990), Rogers et al. (1990), Babb et al. (1991), Blackie et al. (1991), Kohrt et al. (1991), Roca et al. (1992), Kastello et al. (1993), Stevenson et al. (1994), Kasch et al. (1995), McClaran et al. (1995), Rowland and Boyajian (1995), Gore et al. (1996), Podolsky et al. (1996), Proctor and Beck (1996), Rowland and Cunningham (1997), Tanaka et al. (1997), Harms et al. (1998), Marven et al. (1998), Putman et al. (1998), Katayama et al. (1999), Pellegrino et al. (1999), Hunter et al. (2000), Nottin et al. (2000), Loftin et al. (2001), Riddell et al. (2001), Schiller et al. (2001), Poole et al. (2002), Neder et al. (2003), Ocel et al. (2003), Pimentel et al. (2003), Volianitis et al. (2003), Olfert et al. (2004), Petersson et al. (2004), Nourry et al. (2005), Osborne and Schneider (2005), Stickland et al. (2006), and Woo et al. (2006).